

Grassmannian and chiral anomaly

Booss-Bavnbek, Bernhelm; Wojciechowksi, K.P.

Publication date:
1996

Document Version
Publisher's PDF, also known as Version of record

Citation for published version (APA):
Booss-Bavnbek, B., & Wojciechowksi, K. P. (1996). *Grassmannian and chiral anomaly*. Roskilde Universitet. Tekster fra IMFUFA No. 312 <http://milne.ruc.dk/ImfufaTekster/>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact rucforsk@kb.dk providing details, and we will remove access to the work immediately and investigate your claim.

TEKST NR 312

1996

**Grassmannian
and
Chiral Anomaly**

B. Booss-Bavnbek

K. P. Wojciechowski

TEKSTER fra

IMFUFA

ROSKILDE UNIVERSITETSCENTER
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

IMFUFA, Roskilde University, P.O.Box 260, 4000 Roskilde, Denmark
DEPARTMENT OF STUDIES IN MATHEMATICS AND PHYSICS AND THEIR
FUNCTIONS IN EDUCATION, RESEARCH, AND APPLICATIONS

GRASSMANNIAN AND CHIRAL ANOMALY

B. Booss-Bavnbek, K. P. Wojciechowski

IMFUFA text no. 312

26 pages

ISSN 0106-6242

ABSTRACT. We discuss the Grassmannian of self-adjoint global elliptic boundary conditions with γ_5 - and gauge-invariance of the domain for the Dirac operator over the 4-ball coupled to a gauge configuration with non-trivial curvature form arising from local chiral anomaly. We show that this space contains a variety of boundary conditions in addition to the spectral Atiyah-Patodi-Singer projection and that some of them, like the Calderón projector, imply global (strong) chiral symmetry.

We undertake to illuminate questions put to us by the physicists Gianni Morchio and Franco Strocchi (Pisa) and Andrzej Trautman (Warsaw and Trieste) and propose some possible answers to their questions.

GRASSMANNIAN AND CHIRAL ANOMALY *

B. BOOSS-BAVNBEK

*Institut for matematik og fysik
Roskilde University
DK-4000 Roskilde, Denmark
e-mail: booss@mmf.ruc.dk*

and

K. P. WOJCIECHOWSKI

*Dept. Math., IUPUI (Indiana/Purdue)
Indianapolis IN 46202-3216, U.S.A.
e-mail: kwojciechowski@math.iupui.edu*

Abstract. We discuss the Grassmannian of self-adjoint global elliptic boundary conditions with γ_5 - and gauge-invariance of the domain for the Dirac operator over the 4-ball coupled to a gauge configuration with non-trivial curvature form arising from local chiral anomaly. We show that this space contains a variety of boundary conditions besides the spectral Atiyah-Patodi-Singer projection and that some of them, like the Calderón projector, imply global (strong) chiral symmetry.

Key words: Atiyah-Patodi-Singer index theorem, Calderón projector, chiral symmetry, Dirac operator, elliptic boundary problems, γ_5 -symmetry, gauge-invariance, pure gauge, quantum chromodynamics, pseudo-differential Grassmannian.

Introduction

It is implicit in recent literature on quantum field theory that γ_5 - and gauge-invariance of a self-adjoint elliptic boundary condition P for the (massless) Dirac operator \mathcal{D}_A over the 4-ball coupled to a non-trivial gauge configuration A which is pure gauge at the boundary lead to *global* chiral anomaly, i.e. the integers n_+ and n_- of the zero frequency modes of positive and negative chirality can not be expected to be identical and that their difference will depend on the winding number $\deg A$ associated with the non-negligible external field of the gauge configuration and arising from *local* chiral symmetry breaking. The reason for this belief may be an undue prominence of the spectral Atiyah-Patodi-Singer (APS) boundary condition $\Pi_>$ for the corresponding partial Dirac operator \mathcal{D}_A^+ . As observed by M. Ninomiya and C.I. Tan in [20], imposing APS boundary conditions does lead to global (strong) chiral asymmetry. But the APS is far from being the only reasonable elliptic, self-adjoint, γ_5 -invariant, and gauge-invariant boundary condition.

In [17], G. Morchio and F. Strocchi noticed the role played by boundary terms for the discussion of spontaneous symmetry breaking. They saw no reason to question the *local* chiral anomaly, i.e. the existence of the correction term which invalidates the continuity equation for the chiral current (reconstructed through the Matthews-

* Working paper, January 19, 1996, expanding discussions with Gianni Morchio and Franco Strocchi (Pisa), and Andrzej Trautman (Warsaw and Trieste)

Salam formula for the fermion correlation functions and mathematically expressed by the curvature of the underlying connection). Instead, they addressed the *global* implications of this term, essentially the question whether its volume integral vanishes or not. That must depend on the boundary conditions chosen. In that connection they questioned the belief in the necessity of imposing the APS boundary condition leading to global (strong) chiral anomaly and put forward some arguments supporting the view that

- (a) the *ellipticity* and *self-adjointness* of the boundary condition, providing a nicely spaced discrete real spectrum with only accumulation point at $\pm\infty$;
 - (b) the γ_5 -*invariance* of the boundary condition; and
 - (c) the invariance of the domain under any *globally defined gauge* transformation
- might be compatible with
- (d) *global (strong) chiral symmetry*, i.e. the vanishing of $n_+ - n_-$.

The compatibility of the four properties would lead to considerable simplifications in the calculations and a new understanding of the physical effects especially connected with the θ term in quantum chromodynamics. As explained in [17], Section 5.5, chiral symmetry would imply that the exponential

$$e^{\theta \int F^* F dx} = e^{i\theta[(n_+ - n_-)(A)]} = \lim_{m \rightarrow 0} m^{-(n_+ + n_-)} \det \mathcal{D}_A^{m, \theta} =: f_A(\theta)$$

of the effective action would become independent of the fermion angle θ of the fermion mass term, and vice versa. On the mathematical level one could simplify considerably the ζ function regularization of the determinant bundle with

$$\begin{aligned} \zeta'(s) &= - \sum_{\lambda \in \text{spec } \mathcal{D}_A} \ln \lambda e^{s \ln |\lambda|^2} \\ &= - [(n_+ + n_-) \ln m + i(n_+ - n_-)\theta_m] e^{-s \ln m^2} - \sum_{\lambda \neq 0} \ln |\lambda|^2 e^{-s \ln |\lambda|^2}. \end{aligned}$$

It follows that we must discuss the number of the zero modes. We shall show that the Morchio-Strocchi conjecture is provable and that global (strong) chiral symmetry is indeed compatible with the other requirements. It seems, moreover, that there are no gauge-theoretic arguments for selecting the chiral anomaly generating Atiyah-Patodi-Singer boundary conditions.

In Section 1 we give a review of the theory of elliptic boundary value problems for any total Dirac operator \mathcal{D} and, in case of even-dimensional manifolds, for the half Dirac operator \mathcal{D}^+ arising from the chiral decomposition $\mathcal{D} = \begin{pmatrix} 0 & \mathcal{D}^- \\ \mathcal{D}^+ & 0 \end{pmatrix}$.

Our definition of the ellipticity is somewhat new, and we hope that it makes this concept more accessible to non-specialists. The standard definition (see for instance [6], Chapter 18) is clearly a special case of the concept we offer here, but we suspect that actually both notions of the ellipticity of a boundary problem are equivalent. We apply the classical concepts of Cauchy data spaces and of the Calderón projector $\mathcal{P}_+(\mathcal{D}^+)$ in order to investigate the ellipticity condition. Then we review the elliptic boundary problems used by physicists and mathematicians. At the end of

the section we discuss the Grassmannian $\text{Gr}(\mathcal{D}^+)$ of all generalized Atiyah-Patodi-Singer conditions, i.e. the space of all pseudo-differential projections with the same principal symbol as the Calderón projector. It provides a natural space of elliptic boundary conditions for the partial Dirac operator \mathcal{D}^+ .

In Section 2 we present a theory of γ_5 -invariant self-adjoint elliptic boundary problems for any total Dirac operator $\mathcal{D} = \begin{pmatrix} 0 & \mathcal{D}^- \\ \mathcal{D}^+ & 0 \end{pmatrix}$. We show that in a natural way each $P \in \text{Gr}(\mathcal{D}^+)$ defines a projection $P^\# \in \text{Gr}_{\gamma_5}^*(\mathcal{D})$, the Grassmannian of γ_5 -invariant self-adjoint elliptic boundary problems for the total Dirac operator.

In Section 3 we discuss the twisted Dirac operator \mathcal{D}_A over a 4-ball V (of large radius) coupled to a vector potential A (a non-trivial gauge configuration which is pure gauge at the boundary). We fix the notation, especially concerning the auxiliary vector bundles and underlying metrics and connections. We exploit the product form $\mathcal{D}_A^+ = N(\partial_r + \not\partial_B)$ of the twisted Dirac operator near the boundary and investigate the effect of changes of the connection form A onto the corresponding field B over the boundary entering into the definition of the boundary (*tangential*) Dirac operator $\not\partial_B$. We show that not only the spectral Atiyah-Patodi-Singer projection $\Pi_{\geq}(\not\partial_B)$ defines a gauge-invariant element of $\text{Gr}_{\gamma_5}^*(\mathcal{D}_A)$, but also the Calderón projector $\mathcal{P}_+(\mathcal{D}_A^+)$.

Assume that the 4-ball V is equipped with the standard metric which makes (according e.g. to J.R. Schmidt and A.M. Bincer [23]) the boundary Dirac operator invertible and its spectrum symmetric. Assume also that the vector potential A is pure gauge at the boundary. Then one obtains the well-known formula

$$\text{index } \mathcal{D}_{A, \Pi_{\geq}} = n_+(\Pi_{\geq}) - n_-(\Pi_{\geq}) = \deg(A) \neq 0 \quad (1)$$

derived by Ninomiya and Tan [20] from the Atiyah-Patodi-Singer index theorem. The proof will be discussed below.

Replacing the spectral projection $\Pi_{\geq}(\not\partial_B)$ by the Calderón projector $\mathcal{P}_+(\mathcal{D}_A^+)$ we get, contrary to (1)

$$\text{index } \mathcal{D}_{A, \mathcal{P}_+} = n_+(\mathcal{P}_+) - n_-(\mathcal{P}_+) = 0, \quad (2)$$

since, by definition, $n_+(\mathcal{P}_+)$ and $n_-(\mathcal{P}_+)$ vanish. From (2) we get the main result of this paper, namely that any twisting of the Euclidean Dirac operator over the 4-ball by means of a connection in an auxiliary bundle of coefficients can be 'lifted' to a suitable section in the Grassmannian providing global (strong) chiral symmetry:

Theorem 0.1 *Let $\text{Conn}_0(V \times \mathbb{C}^2)$ denote the affine space of smooth connections in the coefficients' bundle $V \times \mathbb{C}^2$ over the 4-ball V which are pure gauge at the boundary. Then there exists a smooth map*

$$\mathcal{R} : \text{Conn}_0(V \times \mathbb{C}^2) \ni A \mapsto \mathcal{R}(A) \in \text{Gr}(\mathcal{D}_A^+)$$

which satisfies (a)-(d).

From a mathematical point of view, the preceding theorem provides the best solution available for the compatibility problem of global (strong) chiral symmetry

with local symmetry breaking when the section \mathcal{R} of the Grassmannian is based on the Calderón projector, since it provides

$$n_{\pm}(\mathcal{R}(A)) = 0 \quad (3)$$

for any connection A .

In Section 4 we discuss various alternative self-adjoint, elliptic, γ_5 - and gauge-invariant boundary problems for the twisted Dirac operator \mathcal{D}_A with vanishing $n_+ - n_-$ to give a clear and complete picture of the variety of possibilities of obtaining compatibility of global (strong) chiral symmetry with local chiral asymmetry.

In Section 5 we make some remarks on the construction of the Calderón projector and its relation to the spectral projection (Atiyah-Patodi-Singer boundary condition).

In the Appendix we determine, based on a result by L. Nicolaescu, the adiabatic limit of the Cauchy data spaces of the twisted Dirac operator for the radius of the 4-ball $R \rightarrow \infty$ in terms of the eigenfunctions of the corresponding tangential Dirac operator over the 3-sphere.

1. A Brief Review of the Theory of Elliptic Boundary Problems for Dirac Operators

Let M be a compact smooth oriented Riemannian manifold with boundary Y , and let $S \rightarrow M$ be a bundle of Clifford modules with compatible Hermitian structure and connection (covariant derivative) D . The (total) Dirac operator

$$\mathcal{D} : C^\infty(M; S) \longrightarrow C^\infty(M; S).$$

is obtained by suitably composing the connection $D : C^\infty(M; S) \rightarrow C^\infty(M; T^*M \otimes S)$ with the Clifford multiplication $c : C^\infty(M; TM \otimes S) \rightarrow C^\infty(M; S)$.

Clearly \mathcal{D} is an elliptic differential operator. It is formally self-adjoint and Green's formula holds for all spinors s and s' :

$$(\mathcal{D}s, s') - (s, \mathcal{D}s') = - \int_Y \langle N(y)(s|_Y), s'|_Y \rangle, \quad (4)$$

where $N := c(\mathbf{n}) : S|_Y \rightarrow S|_Y$ denotes the unitary bundle isomorphism given by Clifford multiplication by the inward unit tangent vector.

To proceed further we assume that M is an even-dimensional manifold. Let γ_5 denote the global section of $\text{Hom}(S, S)$ defined locally by

$$\gamma_5 := c(e_1) \dots c(e_k),$$

where $\{e_\mu\}$ is any positively oriented orthonormal local frame of tangent vectors and k denotes the dimension of the manifold M . Since k is even, S splits into subbundles S^\pm spanned by the eigensections of γ_5 corresponding to the eigenvalue ± 1 , if k is divisible by 4, or $\pm i$ otherwise; the Clifford multiplication N switches between $S^\pm|_Y$ and $S^\mp|_Y$; and the Dirac operator splits correspondingly into components

$$\mathcal{D} = \begin{pmatrix} 0 & \mathcal{D}^- \\ \mathcal{D}^+ & 0 \end{pmatrix}$$

such that the *right partial (half) Dirac operator* $\mathcal{D}^+ : C^\infty(M; S^+) \rightarrow C^\infty(M; S^-)$ is formally adjoint to the *left partial operator* $\mathcal{D}^- : C^\infty(M; S^-) \rightarrow C^\infty(M; S^+)$.

To simplify the exposition we assume that the Riemannian metric and the Hermitian structure are product near the boundary. Let us point out that the results presented here remain true also for non-product structures. Admitting non-product structures, however, makes the analysis more complicated especially when one wants to discuss the ζ regularized determinant and related asymptotic expansions.

Close to the boundary, the total Dirac operator splits into the following *product form*

$$\mathcal{D} = \Gamma(\partial_r + B) = \begin{pmatrix} 0 & -N^{-1} \\ N & 0 \end{pmatrix} \left(\partial_r + \begin{pmatrix} \not\partial & 0 \\ 0 & -N \not\partial N \end{pmatrix} \right), \quad (5)$$

where r denotes the inward oriented normal (radial) coordinate in a collar neighbourhood of the boundary and $\not\partial : C^\infty(Y; S^+) \rightarrow C^\infty(Y; S^+)$ denotes the (*tangential*) Dirac operator over the boundary. Notice that in fact

$$\Gamma^2 = -1 \quad \text{and} \quad \Gamma B = -B\Gamma$$

as required by the formal self-adjointness of \mathcal{D} .

We also discuss the operator \mathcal{D}^+ alone. It has the following form on the collar:

$$\mathcal{D}^+ = N(\partial_r + \not\partial) \quad (6)$$

The Dirac operator on an odd-dimensional manifold has the same form $\Gamma(\partial_r + B)$ on the collar. In this case the total operator \mathcal{D} does not split, but the tangential operator is a Dirac operator on an even-dimensional manifold and has therefore the following form:

$$B = \begin{pmatrix} 0 & B^- \\ B^+ & 0 \end{pmatrix}$$

Now we want to discuss the properties of a Dirac operator over a compact manifold with boundary. In the beginning we shall not distinguish between the cases of even- and odd-dimensional manifolds and whether we treat the total or the half Dirac operator. So, let $\mathcal{A} \in \{\mathcal{D}, \mathcal{D}^+\}$ with $\mathcal{A} : C^\infty(M; E) \rightarrow C^\infty(M; F)$, $E, F \in \{S, S^\pm\}$, and product form $\mathcal{A} = \Gamma(\partial_r + B)$ near the boundary Y .

Contrary to the case of a closed manifold, the space

$$\mathcal{H}(\mathcal{A}, \infty) = \{s \in C^\infty(M; E) \mid \mathcal{A}s = 0\}$$

of solutions of the operator \mathcal{A} is an infinite-dimensional subspace of $C^\infty(M; E)$. There is also a question of regularity of the solutions. Let s denote a solution of \mathcal{A} , which is an element of the space $L^2(M; E)$ (or more generally of $H^k(M; E)$ the k -th Sobolev space). In general, it does not follow that s is a smooth section of S . This leads us to the following general definition of the ellipticity.

Definition 1.1 Let \mathcal{A} be a Dirac operator on M and let $\mathcal{A}_\mathcal{R}$ be a closed extension of \mathcal{A} in $L^2(M, S)$ with domain \mathcal{R} . We call $\mathcal{A}_\mathcal{R}$ an *elliptic boundary problem* for the operator \mathcal{A} , if and only if the following two conditions are satisfied:

- (I) The extension $\mathcal{A}\mathcal{R} : \mathcal{R} \rightarrow L^2$ of \mathcal{A} is a Fredholm operator.
- (II) The spaces $\ker \mathcal{A}\mathcal{R}$ and $\operatorname{coker} \mathcal{A}\mathcal{R}$ are (respectively in the case of the cokernel: can be represented as) finite-dimensional subspaces of the spaces of smooth sections.

Remark 1.2 Let $\mathcal{H}(\mathcal{A})$ denote the space of all L^2 solutions of the operator \mathcal{A} . Then we may reformulate condition (II) in the definition as follows:

$$\mathcal{R} \cap \mathcal{H}(\mathcal{A}) \subset C^\infty \quad \text{and} \quad \mathcal{R}^* \cap \mathcal{H}(\mathcal{A}^*) \subset C^\infty,$$

where \mathcal{R}^* denotes the domain of the adjoint operator.

It seems at first sight that the boundary Y does not appear in the definition, but usually the domain \mathcal{R} is defined by a condition posed on the spinors on the boundary: Let γ_0 denote the restriction map $\gamma_0(s)(y) := s(0, y)$. It gives a continuous map $\gamma_0 : H^1(M; E) \rightarrow L^2(Y; E|_Y)$. The condition which determines \mathcal{R} is usually given in the form:

$$\mathcal{R} = \{s \in H^1(M; E) \mid T(\gamma_0(s)) = 0\},$$

where $T : L^2(Y; E|_Y) \rightarrow L^2(Y; G)$ is a 0-th order pseudo-differential operator. Of course T has to satisfy certain additional assumptions to guarantee fulfilment of conditions (I) and (II) from the definition. We have to introduce the Calderón projection in order to explain those conditions.

The *Calderón projector* $\mathcal{P}_+(\mathcal{A})$ is defined as the (without loss of generality orthogonal) projection of $L^2(Y; E)$ onto the *Cauchy data space*, also called *Hardy space* in Clifford analysis:

$$\mathcal{H}_+(\mathcal{A}) := \overline{\{s|_Y \mid s \in C^\infty(M; E) \text{ and } \mathcal{A}s = 0 \text{ in } M \setminus Y\}}^{L^2(Y; E|_Y)}.$$

We discuss the construction of the Calderón projector in the last section of this paper. Let us only point out that the principal symbol $p_+(y; \zeta)$ of $\mathcal{P}_+(\mathcal{A})$ is equal to the orthogonal projection onto the direct sum of the eigenspaces of the automorphism $b(y; \zeta)$ corresponding to the positive eigenvalues. Here b denotes the principal symbol of the tangential operator \mathcal{B} . Now we are ready to formulate the conditions which the operator T has to satisfy:

Definition 1.3 Let $T : C^\infty(Y; E|_Y) \rightarrow C^\infty(Y; E)$ be a pseudo-differential operator of order 0. We call T an *elliptic boundary condition* for the operator \mathcal{A} , if the following conditions are satisfied:

- (I') For any real r , the extension $T^r : H^r(Y; E|_Y) \rightarrow H^r(Y; E)$ of T has a closed range.
- (II') Let $\sigma(T)$ denote the principal symbol of T . Then

$$\operatorname{range}(\sigma(T)) = \operatorname{range}(\sigma(T) \circ p_+).$$

In particular the restriction $\sigma(T)|_{\operatorname{range}(p_+)} : \operatorname{range}(p_+) \rightarrow \operatorname{range}(\sigma(T))$ is an isomorphism of vector bundles.

Condition (I') implies that \mathcal{N}_T , the orthogonal projection onto the kernel of T , is a pseudo-differential operator (see [6], Proposition 18.11). For the ease of notation we shall denote the kernel of T by the same letter \mathcal{N}_T . Condition (II') implies that the couple $(\mathcal{N}_T, \mathcal{H}_+(\mathcal{A}))$ is a *Fredholm pair of subspaces*, i.e. a pair of closed subspaces with finite-dimensional intersection and with sum of finite codimension (then the difference of these two dimensions is called the *index* of the pair; see [6]). More precisely we have the following result:

Proposition 1.4 *Let T be as in Definition 1.3. Then the couple $(\mathcal{N}_T, \mathcal{H}_+(\mathcal{A}))$ is a Fredholm pair of subspaces in $L^2(Y; E|_Y)$ with*

$$\text{index}(\mathcal{N}_T, \mathcal{H}_+(\mathcal{A})) = \text{index} \{T \circ \mathcal{P}_+(\mathcal{A}) : \mathcal{H}_+(\mathcal{A}) \rightarrow \text{range}(T)\} = \text{index } \mathcal{A}_T,$$

where \mathcal{A}_T denotes the realization of the operator \mathcal{A} with the domain

$$\{s \in H^1(M; E) \mid T(\gamma_0(s)) = 0\}.$$

Proof We show that $\text{index}(\mathcal{N}_T, \mathcal{H}_+(\mathcal{A})) = \text{index} \{T\mathcal{P}_+(\mathcal{A}) : \mathcal{H}_+(\mathcal{A}) \rightarrow \text{range}(T)\}$ and refer to [6], Theorem 20.8 for the proof of the second equality.

Let us assume that z is an element of $\mathcal{N}_T \cap \mathcal{H}_+(\mathcal{A})$. This implies that

$$Tz = 0 \quad \text{and} \quad \mathcal{P}_+(\mathcal{A})z = z, \tag{7}$$

which shows that z is an element of the kernel of the operator $T\mathcal{P}_+(\mathcal{A})$. Let us also observe that the second equality of (7) shows that there exists a uniquely determined s , such that $\mathcal{A}s = 0$ and $\gamma_0(s) = z$. Therefore we have shown:

$$\ker \mathcal{A}_T \cong \ker T\mathcal{P}_+(\mathcal{A}) = \mathcal{N}_T \cap \mathcal{H}_+(\mathcal{A}).$$

Now let us assume that w is an element of $(\mathcal{N}_T + \mathcal{H}_+(\mathcal{A}))^\perp$. It means that w is perpendicular to $\mathcal{H}_+(\mathcal{A})$, hence $\mathcal{P}_+(\mathcal{A})w = 0$ and that w is perpendicular to the kernel of T . Therefore there exists q such that $w = T^*q$, which provides the identification of $\ker \mathcal{P}_+(\mathcal{A})T^*$ with the orthogonal complement of the sum and thus ends the proof of the proposition. \square

Now we review some examples of boundary value problems for Dirac operators. We begin with the theoretically most obvious example:

Example 1.1 We put $T := \mathcal{P}_+(\mathcal{A})$, the *Calderón projector* of \mathcal{A} . This is an elliptic boundary condition and in the case of the total Dirac operator $\mathcal{A} := \mathcal{D}$ it provides us with a self-adjoint, elliptic boundary problem for the operator \mathcal{D} . In the case of $\mathcal{A} = \mathcal{D}^+$ we obtain a closed (unbounded) Fredholm operator $\mathcal{D}_{\mathcal{P}_+(\mathcal{D}^+)}^+$ with index equal to 0.

We have to choose different boundary conditions in order to obtain a non-trivial index.

Example 1.2 We still discuss the operator \mathcal{D}^+ . Then the tangential Dirac operator ∂ is a self-adjoint elliptic differential operator over the closed manifold Y and it has an orthogonal complete system of eigenspinors providing a spectral decomposition of $L^2(Y; S^+|_Y)$. Let $\Pi_{\geq}(\partial)$ and $\Pi_{<}(\partial)$ denote the orthogonal projections

of $L^2(Y; S^+|_Y)$ onto its subspace spanned by the eigenspinors corresponding to the non-negative, respectively negative eigenvalues of \not{D} . Those *spectral projections* are pseudo-differential operators and the principal symbol of $\Pi_{\geq}(\not{D})$ is equal to p_+ . Therefore $\Pi_{\geq}(\not{D})$ is an elliptic boundary condition for the operator \mathcal{D}^+ . The problem $\mathcal{D}^+_{\Pi_{\geq}(\not{D})}$ was studied by Atiyah, Patodi and Singer in [2], where they gave the famous index formula for the operator $\mathcal{D}^+_{\Pi_{\geq}(\not{D})}$. We will discuss this formula later.

Example 1.3 Let us discuss the Atiyah-Patodi-Singer problem for the total operator \mathcal{D} in the case of an *odd-dimensional* manifold M . In this case their index formula gives

$$\text{index } \mathcal{D}_{\Pi_{\geq}(\not{D})} = \dim \ker \not{D}^+,$$

i.e. for non-vanishing kernel of the tangential operator the Atiyah-Patodi-Singer problem for the total, symmetric Dirac operator is not self-adjoint, and its index is not stable under small deformations.

On the other hand, Green's formula shows that in the case $\ker(\not{D}) = \{0\}$ the operator $\mathcal{D}_{\Pi_{\geq}(\not{D})}$ is a self-adjoint operator.

Example 1.4 For odd-dimensional M we have two natural *local elliptic boundary conditions* π_{\pm} for the total Dirac operator defined by the chiral projection of $S|_Y$ onto $(S|_Y)^{\pm}$. We then get $\text{index } \mathcal{D}_{\pi_-} - \text{index } \mathcal{D}_{\pi_+} = \text{index } \not{D}^+$. But $\text{index } \mathcal{D}_{\pi_{\pm}}$ vanishes by Green's formula so that we get the illustrious cobordism theorem from the preceding equality, namely the vanishing of the index of any (half) Dirac operator over a closed even-dimensional manifold Y , if the operator can be written as the (half) tangential operator of a (total) Dirac operator over an odd-dimensional manifold M with $\partial M = Y$.

Example 1.5 Also on any odd-dimensional manifold M we have the *chiral bag model*: Let S be a bundle of Clifford modules and $\mathcal{D} : C^{\infty}(M; S) \rightarrow C^{\infty}(M; S)$ a compatible Dirac operator over M . For any natural n and any smooth map $g : Y \rightarrow U(n)$ we get a self-adjoint elliptic operator \mathcal{A}_g acting like

$$\mathcal{A}_g := \begin{pmatrix} n\mathcal{D} & 0 \\ 0 & -n\mathcal{D} \end{pmatrix}$$

with

$$\text{dom } \mathcal{A}_g := \left\{ \begin{pmatrix} s_1 \\ s_2 \end{pmatrix} \in H^1(M; (S \otimes \mathbb{C}^n) \oplus (S \otimes \mathbb{C}^n)) \mid s_2|_Y = (\Gamma \otimes g)s_1|_Y \right\}$$

where $n\mathcal{D} := \mathcal{D} \otimes \text{Id}_{\mathbb{C}^n}$.

Example 1.6 Now we return to the case of even-dimensional M and replace the spectral projection (i.e. the Atiyah-Patodi-Singer boundary condition for the partial Dirac operator \mathcal{D}^+) by projections belonging to the *Grassmannian* $\text{Gr}(\mathcal{D}^+)$ of generalized Atiyah-Patodi-Singer boundary conditions for \mathcal{D}^+ . The space $\text{Gr}(\mathcal{D}^+)$ is defined as the space of pseudo-differential projections with principal symbol equal to the orthogonal projection p_+ . Here 'projection' means 'idempotent' (i.e. $P = P^2$). The Grassmannian is locally pathwise connected and has countably many connected

components; two projections P_1, P_2 belong to the same component, if and only if the *virtual codimension*

$$i(P_2, P_1) := \text{index} \{P_2 P_1 : \text{range } P_1 \rightarrow \text{range } P_2\} = \text{index} (\text{Id} - P_2, P_1) \quad (8)$$

of P_1 in P_2 vanishes. Here $\text{index} (\text{Id} - P_2, P_1)$ denotes the index of the Fredholm pair (of ranges of the projections). The higher homotopy groups of each connected component are given by Bott periodicity. According to Proposition 1.4 we have $\text{index } \mathcal{D}_P^+ = i(P, \mathcal{P}_+(\mathcal{D}^+))$ for all $P \in \text{Gr}(\mathcal{D}^+)$.

Note: Important elements of $\text{Gr}(\mathcal{D}^+)$ are the *weighted spectral projections* $\Pi_{\geq a}(\not{D})$ with a cut of the spectrum at an arbitrary real a . More precisely, they are defined as the orthogonal projections onto the direct sum of the eigenspaces of the tangential Dirac operator \not{D} for eigenvalues $\geq a$.

Another important example is the *Calderón projector* $\mathcal{P}_+(\mathcal{D}^+)$ discussed in Example 1.1. Note that the Calderón projector is defined in global data whereas the spectral projections are defined by the data of the tangential Dirac operator, i.e. by data that live on the boundary. In any case, they have the same principal symbol p_+ , but in general belong to different connected components of the Grassmannian.

Example 1.7 Let us now assume that M is an odd-dimensional manifold. In this case we have also an important *self-adjoint Grassmannian* $\text{Gr}^*(\mathcal{D})$ of self-adjoint boundary conditions of Atiyah-Patodi-Singer type. This is the subspace of $\text{Gr}(\mathcal{D})$, which consists of those (orthogonal) projections P , which satisfy the condition

$$-\Gamma P \Gamma = \text{Id} - P.$$

Any element of $\text{Gr}^*(\mathcal{D})$ defines a self-adjoint elliptic boundary value problem for the operator \mathcal{D} .

2. γ_5 -Invariant Elliptic Boundary Problems

In this section we discuss the boundary value problems related to the physical situation described in the Introduction. We now assume that the manifold M is even-dimensional. We begin with two typical examples of elliptic boundary problems:

Examples 2.1 (a) The tangential Dirac operator \not{D} is a self-adjoint elliptic differential operator over the closed manifold Y . As in Section 1, $\Pi_{\geq}(\not{D})$ and $\Pi_{<}(\not{D})$ denote the orthogonal projections of $L^2(Y; S^+|_Y)$ onto its subspace spanned by the eigen-spinors corresponding to the non-negative, respectively negative eigenvalues of \not{D} . Choosing

$$\Pi := \begin{pmatrix} \Pi_{\geq}(\not{D}) & 0 \\ 0 & N \Pi_{<}(\not{D}) N^{-1} \end{pmatrix} \quad (9)$$

as boundary condition we obtain an operator \mathcal{D}_Π which is a self-adjoint Fredholm operator with smooth kernel and cokernel and nicely spaced spectrum, such that the invariants

$$\eta_{\mathcal{D}_\Pi}(0), \quad \zeta_{\mathcal{D}_\Pi}(0), \quad \text{and} \quad \zeta'_{\mathcal{D}_\Pi}(0)$$

are defined in exactly the same way as in the closed case. Here the product structure near the boundary is important. Actually it turns out that the spectrum is symmetric due to γ_5 -symmetry, hence the η -invariant vanishes, see Proposition 2.2b below.

(b) If we set (following [14])

$$T := \frac{1}{2} \begin{pmatrix} 1 & N^{-1} \\ N & 1 \end{pmatrix} \quad (10)$$

we obtain a realization \mathcal{D}_T with the same properties as listed for \mathcal{D}_Π in (a).

The reason why the listed properties are in line with each other in these two cases is that $P \in \{\Pi, T\}$ satisfy the same two basic conditions (for details see [5] and [6]):

- The first condition is the *ellipticity* (well-posedness) of the projection P defining the boundary condition explained in Definitions 1.1, 1.3 above.
- The second condition is the *self-adjointness* of the L^2 -extension. As discussed in Example 1.7 this is a symmetry condition in the normal derivatives, more precisely demanding $\text{Id} - P = -\Gamma P \Gamma$.

One decisive difference between the two boundary conditions Π and T defined in equations (9) and (10) lies in the γ_5 -symmetry: In view of the chiral splitting γ_5 takes the form $\gamma_5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$ for dimension of M divisible by 4 (otherwise we multiply by the imaginary unit i), hence $\gamma_5 \Pi \gamma_5 = \Pi$, i.e. Π is γ_5 -invariant, but $\gamma_5 T \gamma_5 = \text{Id} - T$, i.e. T is *not* γ_5 -invariant.

The main result of this section is the following proposition.

Proposition 2.2 (a) *The mapping*

$$\text{Gr}(\mathcal{D}^+) \ni P \mapsto P^\# := \begin{pmatrix} P & 0 \\ 0 & N(\text{Id} - P)N^{-1} \end{pmatrix} \quad (11)$$

provides us with the γ_5 -Grassmannian $\text{Gr}_{\gamma_5}^(\mathcal{D})$ of self-adjoint elliptic (well-posed) boundary conditions for the total Dirac operator \mathcal{D} which are all γ_5 -invariant. In particular we have a natural identification*

$$\pi_0(\text{Gr}_{\gamma_5}^*(\mathcal{D})) \simeq \pi_0(\text{Gr}(\mathcal{D}^+)) \simeq \mathbb{Z}. \quad (12)$$

(b) *For all $P^\# \in \text{Gr}_{\gamma_5}^*(\mathcal{D})$ the L^2 realization $\mathcal{D}_{P^\#}$ has a discrete real spectrum. Each eigenvalue is of finite multiplicity and there are no finite accumulation points. The spectrum is symmetric around the origin of the real axis (hence there is no η function).*

(c) *The null space*

$$\ker \mathcal{D}_{P^\#} := \{s \in H^1(M; S) \mid \mathcal{D}(s) = 0 \text{ and } P^\#(s|_Y) = 0\}$$

consists solely of smooth spinors. It is finite-dimensional and splits naturally into the direct sum of a space of spinors of positive chirality of dimension $n_+(P^\#)$ and a space of spinors of negative chirality of dimension $n_-(P^\#)$ with

$$\text{index } \mathcal{D}_P^+ = n_+(P^\#) - n_-(P^\#), \quad (13)$$

where $P^\#$ and P are related through (11).

Proof (a) and the main proposition of (b) follow at once from the general theory of global elliptic boundary problems for the Dirac operator. To prove that the spectrum of $\mathcal{D}_{P^\#}$ is $\lambda \mapsto -\lambda$ symmetric we consider an eigenspinor $s = \begin{pmatrix} s_+ \\ s_- \end{pmatrix} \in \text{dom } \mathcal{D}_{P^\#}$ with $\mathcal{D}s = \lambda s$. Because of the anti-diagonal form of \mathcal{D} this means

$$\mathcal{D}^- s_- = \lambda s_+ \text{ and } \mathcal{D}^+ s_+ = \lambda s_-.$$

Then $\begin{pmatrix} s_+ \\ -s_- \end{pmatrix}$ also belongs to $\text{dom } \mathcal{D}_{P^\#}$ and is an eigenspinor of $\mathcal{D}_{P^\#}$ with eigenvalue $-\lambda$, since

$$\begin{pmatrix} 0 & \mathcal{D}^- \\ \mathcal{D}^+ & 0 \end{pmatrix} \begin{pmatrix} s_+ \\ -s_- \end{pmatrix} = \begin{pmatrix} -\mathcal{D}^- s_- \\ \mathcal{D}^+ s_+ \end{pmatrix} = \begin{pmatrix} -\lambda s_+ \\ \lambda s_- \end{pmatrix} = -\lambda \begin{pmatrix} s_+ \\ -s_- \end{pmatrix}$$

and, trivially,

$$N(\text{Id} - P)N^{-1}(s_-|_Y) = 0 \implies N(\text{Id} - P)N^{-1}(-s_-|_Y) = 0.$$

To see (c), we notice that by definition

$$\ker \mathcal{D}_{P^\#} = \ker \mathcal{D}_P^+ \oplus \ker \mathcal{D}_{N(\text{Id}-P)N^{-1}}^-$$

with $\dim \ker \mathcal{D}_P^+ = n_+(P^\#)$ and $\dim \ker \mathcal{D}_{N(\text{Id}-P)N^{-1}}^- = n_-(P^\#)$. Since $(\mathcal{D}_P^+)^* = \mathcal{D}_{N(\text{Id}-P)N^{-1}}^-$, it follows that $\text{index } \mathcal{D}_P^+ = n_+(P^\#) - n_-(P^\#)$. \square

We close this section with a discussion of the global (strong) chiral anomaly $n_+(P^\#) - n_-(P^\#)$ for the γ_5 -invariant boundary conditions induced by

- the Calderón projector $\mathcal{P}_+(\mathcal{D}^+)$;
- the spectral projection $\Pi_{\geq}(\emptyset)$; and
- the weighted spectral projections $\Pi_{\geq a}(\emptyset)$ for any real a .

For (c) of the preceding proposition it suffices to determine the index of the corresponding problems for the partial (half) Dirac operator.

Examples 2.3 (a) As noticed before, from the definition of the Calderón projector it is immediate that $\ker \mathcal{D}_{\mathcal{P}_+(\mathcal{D}^+)}^+ = 0$. From Green's formula we get that

$$\mathcal{P}_+(\mathcal{D}^-) = N(\text{Id} - \mathcal{P}_+(\mathcal{D}^+))N^{-1},$$

hence

$$\ker \mathcal{D}_{N(\text{Id}-\mathcal{P}_+(\mathcal{D}^+))N^{-1}}^- = \ker \mathcal{D}_{\mathcal{P}_+(\mathcal{D}^-)}^- = 0,$$

hence the index of the elliptic boundary value problem $\mathcal{D}_{\mathcal{P}_+(\mathcal{D}^+)}^+$ vanishes: There is no global (strong) chiral anomaly when imposing the Calderón projector as boundary condition.

(b) Choosing the spectral projection $\Pi_{\geq}(\not{D})$ as boundary condition we have the Atiyah-Patodi-Singer index theorem which gives

$$\text{index } \mathcal{D}_{\Pi_{\geq}(\not{D})}^+ = \int_M \alpha(x) - \frac{1}{2}(\eta_{\not{D}}(0) + \dim \ker \not{D}). \quad (14)$$

Here $\alpha(x)$ denotes the locally defined *index density* of \mathcal{D}^+ which expresses the local chiral anomaly, and

$$\eta_{\not{D}}(z) := \sum_{\lambda \in \text{spec } \not{D} \setminus \{0\}} \text{sign } \lambda |\lambda|^{-z} = \frac{1}{\Gamma(\frac{z+1}{2})} \int_0^\infty t^{\frac{z-1}{2}} \text{tr}(\not{D} e^{-t \not{D}^2}) dt \quad (15)$$

denotes the η -function of the tangential Dirac operator \not{D} . It is

- (i) well defined through absolute convergence for $\Re(z)$ large;
- (ii) it extends to a meromorphic function in the complex plane with isolated simple poles;
- (iii) its residues are given by a local formula; and
- (iv) it has a finite value at $z = 0$,

(see e.g. Gilkey [10]).

One can not expect global (strong) chiral symmetry for the Atiyah-Patodi-Singer boundary problem; in general, none of the expressions in formula (14) will vanish. For sufficiently elementary operators and under additional assumptions the error terms $\eta_{\not{D}}(0)$ and $\dim \ker \not{D}$ will vanish (especially for symmetric spectrum and invertible tangential operator), and fairly easy expressions for $\int_M \alpha(x)$ are obtainable (see below Section 3).

(c) For any real a we consider the weighted spectral projection $\Pi_{\geq a}(\not{D})$. From the Agranovič-Dynin theorem (see [6], p.207) we get

$$\text{index } \mathcal{D}_{\Pi_{\geq a}(\not{D})}^+ = \text{index } \mathcal{D}_{\Pi_{\geq}(\not{D})}^+ + i(\Pi_{\geq a}(\not{D}), \Pi_{\geq}(\not{D})) \quad (16)$$

with the virtual codimension defined in (8) as error term. For $a \geq 0$ the virtual codimension of $\Pi_{\geq}(\not{D}) = \Pi_{\geq 0}(\not{D})$ in $\Pi_{\geq a}(\not{D})$ becomes $\sum_{0 \leq \lambda < a} \dim E_\lambda$ and for $a < 0$ it becomes $-\sum_{a \leq \lambda < 0} \dim E_\lambda$ where E_λ denotes the eigenspace of the tangential operator \not{D} corresponding to λ . Hence, for suitable choice of a we can obtain global (strong) chiral symmetry

$$\text{index } \mathcal{D}_{\Pi_{\geq a}(\not{D})}^+ = 0,$$

even if the index of the Atiyah-Patodi-Singer problem does not vanish. If its index is $\nu \neq 0$, say $\nu > 0$, the spectral cut a must be chosen in such a way that $\sum_{0 \leq \lambda < a} \dim E_\lambda = \nu$.

To exploit the rich structure of the γ_5 -Grassmannian and to investigate the passing from one connected component (*sector*) to another under change of boundary conditions we shall now apply the preceding theory to a specific 4-dimensional problem of gauge theoretic physics, the problem of global (strong) chiral symmetry in the presence of local (weak) chiral symmetry breaking for gauge-invariant boundary conditions.

3. Twisted Dirac Operators over the 4-Ball

Now we address the physics situation. As manifold M we take a 'volume' V in \mathbf{R}^4 . We think of V as a ball of large radius R . Actually, we are interested in the asymptotic situation with $R \rightarrow \infty$. As the bundle of Clifford modules we take

$$V \times (S \otimes \mathbf{C}^2) = S \otimes \mathbf{C}^2, \quad (17)$$

the Clifford bundle of *Euclidean spinors with coefficients in the trivial bundle* $V \times \mathbf{C}^2$ with Clifford action $c(a) \otimes 1$. As the full Dirac operator \mathcal{D} we take a twisted Dirac operator defined by a connection A for $V \times \mathbf{C}^2$ which is *pure gauge* on the boundary of V .

To make all definitions precise, to fix the notation, and to check the parity and the signs we recall that the (*free*) *Euclidean Dirac operator*

$$\mathcal{D} = \begin{pmatrix} 0 & -\frac{\partial}{\partial \bar{q}} \\ \frac{\partial}{\partial \bar{q}} & 0 \end{pmatrix} : C^\infty(V; S) \rightarrow C^\infty(V; S)$$

is canonically defined over \mathbf{R}^4 with

$$\frac{\partial}{\partial \bar{q}} = i \frac{\partial}{\partial x_1} + j \frac{\partial}{\partial x_2} + k \frac{\partial}{\partial x_3} + \frac{\partial}{\partial x_4}, \quad (18)$$

and

$$\frac{\partial}{\partial q} = -i \frac{\partial}{\partial x_1} - j \frac{\partial}{\partial x_2} - k \frac{\partial}{\partial x_3} + \frac{\partial}{\partial x_4}, \quad (19)$$

where the bundle S of Euclidean spinors in 4 dimensions splits into a pair of quaternions $S = V \times (\mathbf{H} \oplus \mathbf{H})$ with Clifford multiplication $c : \text{Cl}_4 \rightarrow \text{Hom}_{\mathbf{C}}(S, S)$ given by the four complex 4×4 matrices

$$c(e_\mu) = \gamma_\mu = \begin{pmatrix} 0 & \sigma_\mu \\ \sigma_\mu & 0 \end{pmatrix} \quad \text{for } \mu = 1, 2, 3 \quad \text{and} \quad c(e_4) = \gamma_4 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$


with $\{\sigma_\mu\}$ denoting the Pauli matrices and $\{e_1, \dots, e_4\}$ a basis of \mathbf{R}^4 . The connection defining the Euclidean Dirac operator is just the standard connection d for S .

Then any connection A for the trivial bundle $V \times \mathbf{C}^2$ defines in a natural way a twisted Dirac operator $\mathcal{D}_A = \mathcal{D} \otimes_A \text{Id}_{\mathbf{C}^2}$. It is characterized by the property

$$\mathcal{D}_A(s \otimes f)(x) = (\mathcal{D}s) \otimes f(x)$$

whenever $(Af)(x) = 0$. It is a true (total) Dirac operator with regard to the induced Clifford multiplication $c(a) \otimes 1$ and the induced connection $d \otimes A$.

Now we must discuss the choice of the connection A . From a physical point of view it does not suffice to consider only the trivial choice, namely the standard connection d in $V \times \mathbf{C}^2$ given by exterior differentiation $\sum_\mu \alpha_\mu e_\mu \mapsto \sum_\mu \alpha_\mu \partial_\mu$. Roughly speaking, the standard connection would correspond to the description of two non-interacting fermions. When these fermions are not considered independently, their interaction on the surface ∂V of the volume V can be described in a canonical

Fig. 1. The collar $[0, \varepsilon) \times \partial V$

way, but expressed in different ways, although it is very much the same thing over the contractible n -ball V (for details see e.g. B.A. Dubrovin, A.T. Fomenko, and S.P. Novikov [7]): One way is to introduce a smooth family h of $SU(2)$ matrices parametrized over ∂V ; this is equivalent to introducing a smooth connection ∇ over the whole ball which is pure gauge at the boundary with regard to h ; or, equivalently, to introduce a vector-valued field $\{A_\mu\}$ which is pure gauge at the boundary with regard to h providing $\nabla_{e_\mu} f = \frac{\partial f}{\partial x_\mu} + A_\mu f - f A_\mu$ for any $f \in C^\infty(V; V \times \mathbb{C}^2)$ and e_μ the μ th basis vector in \mathbb{R}^4 .

The geometrical idea behind demanding the interaction boundary term to behave as pure gauge is to get a non-trivial curvature form $\Omega_\nabla = \sum_{\mu < \nu} F_{\mu\nu} dx_\mu \wedge dx_\nu$ corresponding to an action $\int F_{\mu\nu}^2 dx < \infty$ with

$$\begin{aligned} F_{\mu\nu} &:= [\nabla_{e_\mu}, \nabla_{e_\nu}] = \nabla_{e_\mu} \nabla_{e_\nu} - \nabla_{e_\nu} \nabla_{e_\mu} \\ &= \partial_\mu A_\nu - \partial_\nu A_\mu + [A_\mu, A_\nu] \in L^2 \end{aligned}$$

and $F_{\mu\nu}(x) \rightarrow 0$ for $|x| \rightarrow \infty$. This is equivalent to writing the connection $\nabla = d + \sum_\mu A_\mu$ with $A_\mu(x) \rightarrow h \partial_\mu h^{-1}$ for $|x| \rightarrow \infty$.

The different mathematical descriptions are formalized in the following definition.

Definition 3.1 Let $\text{Conn}(V \times \mathbb{C}^2)$ denote the affine space of smooth connections for the bundle $V \times \mathbb{C}^2$. A connection $A \in \text{Conn}(V \times \mathbb{C}^2)$ is called *pure gauge* at the boundary $\partial V = S^3$, if in a collar $[0, \varepsilon) \times \partial V$ (see Figure 3):

- A does not depend on the normal (radial) coordinate r ; and
- there exists a smooth $h : \partial V \sim S^3 \rightarrow SU(2) \sim S^3$ such that $A = h \circ d \circ h^{-1}$ in the collar, where d denotes the standard connection given by exterior differentiation.

We write $\text{Conn}_0(V \times \mathbb{C}^2)$ for the subspace of connections which are pure gauge at the boundary.

Proposition 3.2 (a) A connection A for $V \times \mathbb{C}^2$ is pure gauge at the boundary if it can be written in the form $A = d - (dh)h^{-1}$ in a collar of the boundary.

(b) If A is pure gauge at the boundary, then the tangential Dirac operator B over ∂V corresponding to the partial Dirac operator \mathcal{D}_A^+ over V takes the form

$$B = \mathcal{D}_{S^3} \otimes_{-(dh)h^{-1}} \text{Id} = (\text{Id} \otimes h) (\mathcal{D} \otimes \text{Id}_{\mathbb{C}^2}) (\text{Id} \otimes h^{-1}) \quad (20)$$

with $\mathcal{D} \otimes \text{Id}_{\mathbb{C}^2} = \mathcal{D} \oplus \mathcal{D}$. Here $\mathcal{D} = \mathcal{D}_{S^3}$ denotes the tangential operator over S^3 corresponding to the Euclidean Dirac operator \mathcal{D}^+ .

Proof To prove (a) we find

$$Af = (hdh^{-1})f = hd(h^{-1}f) = h(h^{-1}df + d(h^{-1})f) = df - (dh)h^{-1}f \quad (21)$$

since $d(hh^{-1})$ vanishes.

To prove (b) we notice that the restriction of A to the boundary takes the form $-(dh)h^{-1}$, therefore we get such a simple form for lifting \not{D} to the auxiliary bundle. For details of the calculation see e.g. [21] and [30]. \square

We shall discuss the Atiyah-Patodi-Singer index formula for the operator \not{D}_A

$$\text{index } (\not{D}_A^+)_{\Pi_2} = \int \alpha(x) - \frac{1}{2} (\eta_B(0) + \dim \ker B).$$

From the preceding proposition we have

$$\eta_B(0) = 2\eta_{\not{D}_{S^3}}(0) \quad \text{and} \quad \dim \ker B = 2 \dim \ker \not{D}_{S^3}.$$

We find $\deg(h)$ for the value of the integral of the index density. This result is actually independent of the choice of the metric. Then

$$\text{index } (\not{D}_A^+)_{\Pi_2} = \deg(h) - \eta_{\not{D}_{S^3}}(0) - \dim \ker \not{D}_{S^3}. \quad (22)$$

The two numbers on the right were found to vanish for the standard metric of \mathbb{R}^4 , slightly modified close to ∂V in a calculation done by J.R. Schmidt and A.M. Bincer in [23], see also Schmidt [22]. In that metric the tangential Dirac operator on the 3-sphere \not{D}_{S^3} has a spectrum symmetric about $\lambda = 0$ and is invertible. More precisely:

Lemma 3.3 [J. R. Schmidt and A. M. Bincer, 1987] *The tangential Dirac operator \not{D} over ∂V corresponding to the (free) Euclidean Dirac operator over a ball $V \subset \mathbb{R}^4$ of radius R in a spherical metric has eigenvalues λ with multiplicity $M(\lambda)$ expressed as*

$$\lambda R = \pm(\frac{1}{2} + \kappa), \quad M(\lambda) = \kappa(\kappa + 1), \quad \kappa = 1, 2, \dots \quad (23)$$

Proof To explain the metric chosen in [23] we must repeat parts of the proof. First rewrite the operators $\not{D}^+ = \frac{\partial}{\partial \bar{q}}$ and $\not{D}^- = -\frac{\partial}{\partial q}$ defined above in (18) and (19) as 2×2 matrices

$$\frac{\partial}{\partial \bar{q}} = \begin{pmatrix} \partial_4 + i\partial_3 & \partial_2 + i\partial_1 \\ -\partial_2 + i\partial_1 & \partial_4 - i\partial_3 \end{pmatrix}$$

and

$$\frac{\partial}{\partial q} = \begin{pmatrix} \partial_4 - i\partial_3 & -\partial_2 - i\partial_1 \\ \partial_2 - i\partial_1 & \partial_4 + i\partial_3 \end{pmatrix}.$$

Then parametrize V by

$$\begin{aligned} x_1 &= r \sin \theta \sin \varphi_1, & x_2 &= r \sin \theta \cos \varphi_1, \\ x_3 &= r \cos \theta \sin \varphi_2, & x_4 &= r \cos \theta \cos \varphi_2, \\ 0 &\leq \varphi_{1,2} \leq 2\pi, & 0 &\leq \theta \leq \frac{\pi}{2}, \quad 0 \leq r \leq R \end{aligned}$$

and notice

$$\begin{aligned}\partial_4 \pm i\partial_3 &= e^{\pm\varphi_2} \left(\cos\theta \partial_r - \frac{\sin\theta}{r} \partial_\theta \pm \frac{i}{r \cos\theta} \partial_{\varphi_2} \right), \\ \partial_2 \pm i\partial_1 &= e^{\pm\varphi_1} \left(\sin\theta \partial_r + \frac{\cos\theta}{r} \partial_\theta \pm \frac{i}{r \sin\theta} \partial_{\varphi_1} \right).\end{aligned}$$

To cast

$$\mathcal{D} = \begin{pmatrix} 0 & -\frac{\partial}{\partial \bar{q}} \\ \frac{\partial}{\partial \bar{q}} & 0 \end{pmatrix}$$

into the required product form

$$\mathcal{D} = \Gamma(\partial_r + \mathcal{B})$$

close to the boundary, the partial Dirac operators $\frac{\partial}{\partial \bar{q}}$ and $\frac{\partial}{\partial q}$ are replaced by

$$\widetilde{\mathcal{D}}^+ := Q \frac{\partial}{\partial \bar{q}} R^{-1}, \quad \widetilde{\mathcal{D}}^- = -R \frac{\partial}{\partial q} Q^{-1},$$

with

$$\begin{aligned}R &:= (r^3 \sin\theta \cos\theta)^{\frac{1}{2}} \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} e^{-i\varphi_-} & 0 \\ 0 & e^{i\varphi_-} \end{pmatrix}, \\ Q &:= (r^3 \sin\theta \cos\theta)^{\frac{1}{2}} \begin{pmatrix} e^{-i\varphi_+} & 0 \\ 0 & e^{i\varphi_+} \end{pmatrix}, \\ \varphi_\pm &:= \frac{1}{2}(\varphi_1 \pm \varphi_2).\end{aligned}$$

This transformation is equivalent to a slight modification of the metric of V near the boundary. One obtains

$$\widetilde{\mathcal{D}}^+ = \partial_r + \mathcal{D}, \quad \widetilde{\mathcal{D}}^- = -\partial_r + \mathcal{D},$$

with

$$\mathcal{D} = \frac{1}{r} \begin{pmatrix} \frac{1}{2} + i\partial_{\varphi_1} + i\partial_{\varphi_2} & \partial_\theta + i \cot\theta \partial_{\varphi_1} - i \tan\theta \partial_{\varphi_2} \\ -\partial_\theta + i \cot\theta \partial_{\varphi_1} - i \tan\theta \partial_{\varphi_2} & \frac{1}{2} - i\partial_{\varphi_1} - i\partial_{\varphi_2} \end{pmatrix}. \quad (24)$$

Setting $r := R$ in (24) we get an explicit form of the tangential Dirac operator. For details of the eigenvalue determination of \mathcal{D} we refer to [23], 3996-3997. \square

Remarks 3.4 (a) We also refer to T. Kori [15] which ensures the same result, namely a symmetric spectrum, not containing zero, for a particular metric set-up coming from a metric over the 4-ball which is product near the boundary. Actually, for Kori's metric the Calderón projector \mathcal{P}_+ and the Atiyah-Patodi-Singer projection Π_+ coincide. Clearly the Calderón projector and the Atiyah-Patodi-Singer projection coincide for the standard metric in two dimensions, but not in four dimensions; see also Section 5 and the Appendix below.

(b) Also from Hitchin [12] it follows directly that the tangential Dirac operator over the 3-sphere in standard metric is non-singular with symmetric spectrum. Explicit calculations of its eigenfunctions were given by Sudbery [27] and Sommen [26].

Recall that $\text{Conn}_0(V \times \mathbb{C}^2)$ denotes the subspace of connections which are pure gauge at the boundary.

Proposition 3.5 [M. Ninomiya and C.I. Tan, 1985] *For a suitable metric we have*

$$\text{index } (\mathcal{D}_A^+)_{\Pi_{\geq}} = \deg(h) \quad (25)$$

for any $A \in \text{Conn}_0(V \times \mathbb{C}^2)$ with $A = h \circ d \circ h^{-1}$ in a collar of ∂V and suitable smooth $h : \partial V \rightarrow \text{SU}(2)$.

Proof The proposition follows at once from the Atiyah-Patodi-Singer Index Theorem (14) and Lemma 3.3. \square

In the same article [20] Ninomiya and Tan pointed out that the Atiyah-Patodi-Singer boundary condition is *natural* or *physical* in the following sense:

Corollary 3.6 *Let*

$$\begin{aligned} \text{Conn}_0(V \times \mathbb{C}^2) \ni A &\mapsto \Pi_h := \Pi_{\geq}(\not{D} \otimes_h \text{Id}_{\mathbb{C}^2}) \in \text{Gr}(\not{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2}) \\ &\mapsto \Pi(h) := (\Pi_h)^{\#} \in \text{Gr}_{\gamma_5}^*(\not{D} \otimes_A \text{Id}_{\mathbb{C}^2}) \end{aligned}$$

denote the mapping provided by the Atiyah-Patodi-Singer boundary condition

$$\Pi(h) := \begin{pmatrix} \Pi_{\geq}(\not{D} \otimes_h \text{Id}_{\mathbb{C}^2}) & 0 \\ 0 & N\Pi_{<}(\not{D} \otimes_h \text{Id}_{\mathbb{C}^2})N^{-1} \end{pmatrix}, \quad (26)$$

where $h : \partial V \rightarrow \text{SU}(2)$ denotes the mapping corresponding to the connection A which is supposed to be pure gauge at the boundary. Then the family of operators $\{\mathcal{D}_{A, \Pi(h)}\}_{A \in \text{Conn}_0(V \times \mathbb{C}^2)}$ which act like $\not{D} \otimes_A \text{Id}_{\mathbb{C}^2}$ with

$\text{dom } \mathcal{D}_{A, \Pi(h)}$

$$:= \left\{ s \in H^1(V; S \otimes \mathbb{C}^2) \mid \Pi(h)(s) = \begin{pmatrix} \Pi_h & 0 \\ 0 & N(\text{Id} - \Pi_h)N^{-1} \end{pmatrix} \begin{pmatrix} s_+ \\ s_- \end{pmatrix} = 0 \right\}$$

satisfies the following three fundamental conditions:

- (a) $\mathcal{D}_{A, \Pi(h)}$ is self-adjoint;
- (b) $\Pi(h)$ is γ_5 -invariant; and
- (c) the domain $\text{dom } \mathcal{D}_{A, \Pi(h)}$ is gauge-invariant.

We recall the meaning of (c): Let $U : V \rightarrow \text{SU}(2)$ define the gauge transformations

$$\begin{aligned} f(x) &\mapsto U(x)f(x)U^{-1}(x), \\ A_{\mu}(x) &\mapsto U(x)A_{\mu}(x)U^{-1}(x) - \frac{\partial U}{\partial x_{\mu}}|_x U^{-1}(x), \end{aligned}$$

then the connection A transforms as follows:

$$\begin{aligned} A_{e_{\mu}}f|_x &\mapsto U(x)(A_{e_{\mu}}f|_x)U^{-1}(x), \\ A &\mapsto UAU^{-1}, \text{ and} \\ \Omega_A &\mapsto U\Omega_A U^{-1} - (dU)U^{-1}. \end{aligned}$$

This motivates the following definition:

Definition 3.7 A smooth family

$$\text{Conn}(V \times \mathbb{C}^2) \ni A \mapsto P(A) \in \text{Gr}_{\gamma_0}^*(\mathcal{P}_A) \cong \text{Gr}(\mathcal{P}_A^*)$$

is *gauge-invariant*, if we have

$$P(A_1) = U^\# P(A) (U^\#)^{-1} \quad (27)$$

for all $A, A' \in \text{Conn}(V \times \mathbb{C}^2)$ where $A_1 := UAU^{-1}$ with arbitrary $U : V \rightarrow \text{SU}(2)$ and $U^\# := \text{Id} \otimes_{\mathbb{C}^2} U$.

Remarks 3.8 (1) Clearly, for any $A, A_1 \in \text{Conn}(V \times \mathbb{C}^2)$ with $A_1 = UAU^{-1}$ we have pointwise (see Palais [21])

$$\mathcal{P}_{A_1} = \mathcal{P} \otimes_{A_1} \text{Id} = U^\# (\mathcal{P} \otimes_A \text{Id}) (U^\#)^{-1} = U^\# \mathcal{P}_A (U^\#)^{-1}.$$

Therefore property (c) - gauge-invariance as defined in (27) - means

$$\mathcal{P}_{A_1, P(A_1)} = U^\# \mathcal{P}_{A, P(A)} (U^\#)^{-1}$$

and, especially,

$$\text{dom}(\mathcal{P}_{A_1, P(A_1)}) = U^\# (\text{dom} \mathcal{P}_{A, P(A)}); \quad (28)$$

i.e. we require that the boundary condition transforms in a correct way under variation of the background operator resp. of the connection.

(2) The change of the number

$$\xi(\mathcal{P}_A; s) := \frac{1}{2} (\eta_{\mathcal{P}_A}(s) + \dim \ker \mathcal{P}_A)$$

for arbitrary change of the connection A , here considered solely as a connection over the closed manifold ∂V , was addressed already in [2] and further investigated in [30]. It turns out that

$$\xi(\mathcal{P}_{A_1}; 0) - \xi(\mathcal{P}_A; 0) \equiv \text{index} \{\mathcal{P}_{A_t}\}_{t \in [0,1]} \pmod{\mathbb{Z}}.$$

Here $t \mapsto A_t$ is a smooth family of connections connecting

$$A = A_0 \quad \text{with} \quad A_1 = UAU^{-1}$$

and $\{\mathcal{P}_{A_t}\}_{t \in [0,1]}$ denotes the corresponding family of Dirac operators parametrized over S^1 or, put differently, the elliptic differential operator of first order over the torus $S^1 \times \partial V$ defined by that family.

Proof of the Corollary

The first two properties are obvious from the choices. The gauge invariance follows from the corresponding transformation law for the tangential operator

$$\mathcal{P} \otimes_{h_1} \text{Id}_{\mathbb{C}^2} = U^\# (\mathcal{P} \otimes_h \text{Id}_{\mathbb{C}^2}) (U^\#)^{-1}, \quad (29)$$

where the smooth families $h, h_1 := U|_{\partial V} h (U|_{\partial V})^{-1} : \partial V \rightarrow \text{SU}(2)$ correspond to the connections A, A_1 which are supposed to be pure gauge at the boundary. Equation (29) implies that the eigenvalues do not change under gauge transformation and that the eigenspaces transform like $E_\lambda(\mathcal{P}_{h_1}) = U E_\lambda(\mathcal{P}_h)$. Hence $\Pi(h)$ satisfies (a), (b), and (c). \square

Note: 1. One should keep in mind that from a geometrical point of view there is no particular reason to choose the Atiyah-Patodi-Singer boundary conditions among all the other boundary problems which, as we shall see, equally satisfy (a)-(c); in other words, it is hard to see what kind of special gauge invariance of the domain should be required to select that particular boundary condition.

2. Moreover, for our purpose it is an unfortunate aspect of the prominent Atiyah-Patodi-Singer boundary condition that its index

$$\text{index } \mathcal{D}_{A, \Pi_h}^+ = n_+(\Pi_h) - n_-(\Pi_h)$$

does not vanish as seen above in Proposition 3.5 stating $\text{index } \mathcal{D}_{\Pi(h)}^+ = \deg(h)$ under suitable conditions about the metric. But there are other quite natural boundary conditions \mathcal{R}_A instead of Π_A which fulfil the conditions (a)-(c) of the Corollary and additionally provide global (strong) chiral symmetry, namely

$$(d) \text{ the vanishing of index } \mathcal{D}_{A, \mathcal{R}_A}^+ = n_+(\mathcal{R}_A) - n_-(\mathcal{R}_A).$$

As announced in the Introduction:

Theorem 3.9 *There exists a smooth map*

$$\mathcal{R} : \text{Conn}_0(V \times \mathbb{C}^2) \ni A \mapsto \mathcal{R}_A \in \text{Gr}(\mathcal{D}_A^+)$$

which satisfies (a)-(d).

Proof It is immediate that

$$A \mapsto \mathcal{R}_A := \mathcal{P}_+(\mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2})$$

satisfies all conditions, where $\mathcal{P}_+(\mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2})$ denotes the Calderón projector of the partial Dirac operator $\mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2}$. \square

Note: 1. A nice feature of the Calderón projector is that it is by definition invariant under parity, i.e.

$$N \circ (\text{Id} - \mathcal{P}_+(\mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2})) \circ N^{-1} = \mathcal{P}_+(\mathcal{D}^- \otimes_A \text{Id}_{\mathbb{C}^2}),$$

whereas the Atiyah-Patodi-Singer boundary condition is invariant under parity only if the tangential operator is invertible.

2. One must expect the existence of a lot of sections of the Grassmannian which lead to families satisfying (a)-(d) (see also Section 4). From a theoretical point of view, the Calderón family is the best solution available for the global (strong) chiral symmetry problems, since it does not only give global (strong) chiral symmetry but also the following proposition. It simplifies radically e.g. the calculation of $\zeta'(s)$ mentioned in the Introduction.

Proposition 3.10 *For the Calderón family the dimensions of the zero frequency modes of positive and negative chirality vanish; i.e. we have*

$$n_{\pm}(\mathcal{P}_+(\mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2})) = 0.$$

4. Alternatives and Further Ramifications

There are various ways of obtaining global (strong) chiral symmetry by imposing elliptic, self-adjoint, γ_5 -symmetric, and gauge-invariant boundary conditions in the presence of local (weak) chiral anomaly (i.e. non-vanishing $\deg h$ for connections which are pure gauge at the boundary). One way is the Calderón projector. It removes all solutions such that kernel and cokernel become trivial. This makes many calculations easy. But since the Calderón projector depends on the gauge configuration also inside the region and not only on the boundary, this must have consequences in, for example, the derivation of identities by variation of the gauge field configuration in a subregion.¹

Instead of removing all solutions by imposing the Calderón projector one can add further solutions to the original Dirac equation with APS boundary condition until one gets global (strong) chiral symmetry. There are three ways to do that.

Let us begin with a given connection A in the auxiliary bundle $V \times \mathbb{C}^2$ which is pure gauge at the boundary. Hence it can be expressed in a collar of the boundary by a mapping $h : S^3 \rightarrow SU(2)$ which has a degree (*topological number*) $\deg h$. If $\deg h = k$ is non-trivial, the dimensions n_+ and n_- of the zero modes (subject to the APS boundary condition Π_h) do not coincide as seen in Proposition 3.5. Then, to get global (strong) chiral symmetry we enlarge the solution spaces until n_+ and n_- become equal. More precisely:

Alternative 4.1 The easiest, but physically hardly very meaningful way of doing the equalization of the solution spaces is to take a second copy of the coefficients bundle $V \times \mathbb{C}^2$ and to choose a connection A' which is pure gauge at the boundary ∂V with a unitary mapping g of opposite degree $-k$.

Then, instead of tensoring the original Euclidean Dirac operator \mathcal{D}^+ solely with the h -connection, we do two twistings: first with h , then with g . The resulting twisted Dirac operator

$$\mathcal{D}'^+ := \mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2} \otimes_{A'} \text{Id}_{\mathbb{C}^2} = \mathcal{D}_A^+ \otimes_{A'} \text{Id}_{\mathbb{C}^2}$$

with coefficients in $\mathbb{C}^2 \otimes \mathbb{C}^2 = \mathbb{C}^4$ admits again an APS boundary condition Π' which is gauge invariant such that

$$\begin{aligned} n_+ - n_- &= \text{index } \mathcal{D}'^+_{\Pi'} = \deg(h \otimes g) \\ &= \deg hg = \deg h + \deg g = k - k = 0. \end{aligned}$$

To get global (strong) chiral symmetry one can also apply a less trivial mirror process:

Alternative 4.2 Instead of twisting the global Dirac operator \mathcal{D}_A^+ over the full 4-ball V it suffices to twist the transversal (tangential) Dirac operator B_h with a connection of opposite degree over the 3-sphere. We get a new operator B'_h . Then we apply the APS spectral projection Π'_h corresponding to the twisted operator B'_h to the original operator \mathcal{D}_A^+ . It follows that Π'_h is an admissible boundary value problem

¹ These calculations will be presented separately.

for \mathcal{D}_A^+ . It belongs to the same Grassmannian as the standard APS projection Π_h and all the nice properties (a)-(c) are guaranteed, but Π'_h belongs to a different connected component. In fact, the index jumps by the winding number yielding global (strong) chiral symmetry.

To see this we recall from equation (20) the formula

$$\Pi_h = (\text{Id} \otimes h)(\Pi_{\geq}(\emptyset) \otimes \text{Id})(\text{Id} \otimes h^{-1}) \quad (30)$$

where $\Pi_{\geq}(\emptyset)$ denotes the APS spectral projection belonging to \mathcal{D}^+ . We recall $\text{index } \mathcal{D}_{A, \Pi_h}^+ = n_+(APS) - n_-(APS) = \deg h$. Here one can prove that Π_h belongs to the same Grassmannian as $\Pi_{\geq}(\emptyset)$ (more precisely as $\Pi_{\geq}(\emptyset) \otimes \text{Id}$), but with the virtual codimension $i(\Pi_h, \Pi_{\geq}(\emptyset)) = \deg h$.

Then we go one tensoring further, namely from B_h to

$$B'_h := (\text{Id} \otimes g)(B_h \otimes \text{Id})(\text{Id} \otimes g^{-1})$$

with $\deg g = -\deg h$. We generalize formula (30) and get a similar formula expressing Π'_h , the APS projection belonging to the operator B'_h , in terms of Π_h , and a new jump formula for the virtual index:

$$\begin{aligned} i(\Pi'_h, \Pi_h) &= i(\text{Id} \otimes g)(\Pi_h \otimes \text{Id})(\text{Id} \otimes g^{-1}), (\Pi_h \otimes \text{Id}) \\ &= \text{index} ((\text{Id} - (\Pi_h \otimes \text{Id}) \oplus \text{Id} \otimes g)(\Pi_h \otimes \text{Id})(\text{Id} \otimes g^{-1}) \circ (\Pi_h \otimes \text{Id})) \\ &= \deg g, \end{aligned}$$

hence

$$\begin{aligned} \text{index } \mathcal{D}_{A, \Pi'_h} &= i(\Pi_{\geq}(B_h), \mathcal{P}_+(\mathcal{D}_A^+)) + i(\Pi_{\geq}(B'_h), \Pi(\mathcal{D}_A^+)) \\ &= \deg h + \deg g = 0. \end{aligned}$$

Note: An attractive feature of Alternative 4.2 is that in fact the (non-free) operator \mathcal{D}_A is not changed; only the boundary condition is changed.

Alternative 4.3 One more alternative is provided by a suitable spectral cut (weighted spectral projection), see above Example 2.3c and, more generally for the problem of uniform choice of the spectral cut, Melrose and Piazza [16].

5. Some Remarks on the Calderón Projector

We begin with the construction of the Calderón projection. Let M be a compact smooth oriented Riemannian manifold with boundary like in Section 1. Let $\tilde{M} = M \cup_Y (-M)$ denote the double of M and $\tilde{S}^+ = S^+ \cup_N S^-$ the corresponding spinor bundle over \tilde{M} . We denote by $\tilde{\mathcal{D}}^+$ the invertible double of the operator \mathcal{D}^+ on \tilde{M} . This is an invertible Dirac operator on \tilde{M} extending \mathcal{D}^+ . The invertibility means that there exists an elliptic pseudo-differential operator \mathcal{G} of order -1 , such that

$$\tilde{\mathcal{D}}^+ \mathcal{G} = \text{Id} \quad \text{and} \quad \mathcal{G} \tilde{\mathcal{D}}^+ = \text{Id}.$$

For any $f \in C^\infty(Y; S^+|_Y)$ we denote by $\delta \otimes f$ the distribution:

$$\langle \delta \otimes f; s \rangle := \int_Y (f; \gamma_0(s)) dy \quad \text{for } s \in C^\infty(\widetilde{M}; \widetilde{S}^+).$$

In fact, the map $f \rightarrow \delta \otimes f$ is the adjoint map to the map γ_0 . Given $f \in C^\infty(Y; S^+|_Y)$ we denote by $F(f)$ the distribution over \widetilde{M} defined as

$$F(f) := \mathcal{G}(\delta \otimes \Gamma f).$$

Now we can give the formula for the Calderón projection:

$$\mathcal{P}_+(\mathcal{D}^+)f := \lim_{u \rightarrow 0} F(f)|_{u \times Y} = \gamma_0 F(f).$$

Though this formula is abstract one can see that basically it depends only on the Green's function of the operator $\widetilde{\mathcal{D}}^+$. The definition extends to $f \in L^2(Y; S^+|_Y)$ by continuity defining a pseudo-differential operator and yielding the Cauchy data space $\mathcal{H}_+(\mathcal{D}^+)$ for the range of $\mathcal{P}_+(\mathcal{D}^+)$, and it turns out that $L^2(Y; S^+|_Y)$ splits into the direct sum

$$L^2(Y; S^+|_Y) = \mathcal{H}_+(\mathcal{D}^+) \oplus N(\mathcal{H}_+(\mathcal{D}^-)),$$

where N denotes Clifford multiplication by the inward unit tangent vector (as above), $\mathcal{D}^- = (\mathcal{D}^+)^*$, and $N(\mathcal{H}_+(\mathcal{D}^-)) = \mathcal{H}_-(\widetilde{\mathcal{D}})$, the outer or right Cauchy data space of the invertible double $\widetilde{\mathcal{D}}^+$.

Next we want to explain the relation of the Calderón projection to the spectral projection of the tangential operator.

Proposition 5.1 *Let \mathcal{D}_A^+ denote the Euclidean Dirac operator over the 4-ball twisted by a connection A which is pure gauge at the boundary with $\deg h$ different than 0. Then its Calderón projection $\mathcal{P}_+(\mathcal{D}_A^+)$ and the spectral projection $\Pi_{>0}(\mathcal{D}_h)$ of the corresponding tangential ('spherical') Dirac operator \mathcal{D}_h belong to different components of the Grassmannian.*

Proof Let $\mathcal{D}^+ \otimes \text{Id}_{\mathbb{C}^2}$ denote the untwisted operator. The operator $\mathcal{D}_A^+ = \mathcal{D}^+ \otimes_A \text{Id}_{\mathbb{C}^2}$ has the same principal symbol, hence $t\mathcal{D}_A^+ + (1-t)(\mathcal{D}^+ \otimes \text{Id}_{\mathbb{C}^2})$ is a path of Dirac operators. It follows from the construction of the Calderón projection that it changes in a continuous way (see Nicolaescu [19] for details). Therefore $\mathcal{P}_+(\mathcal{D}_A^+)$ and $\mathcal{P}_+(\mathcal{D}^+ \otimes \text{Id}_{\mathbb{C}^2})$ belong to the same connected component of the Grassmannian.

The index of $(\mathcal{D}^+ \otimes \text{Id}_{\mathbb{C}^2})_{\Pi_{>0}}$ is equal to 0 (the standard connection has degree 0). This index is equal to $i(\Pi_{>0}(\mathcal{D} \otimes \text{Id}), \mathcal{P}_+(\mathcal{D}^+ \otimes \text{Id}_{\mathbb{C}^2}))$, hence those projections are in the same connected component of the Grassmannian. On the other hand:

$$\deg h = i(\Pi_{>0}(\mathcal{D} \otimes_h \text{Id}), \mathcal{P}_+(\mathcal{D}_A^+)) \neq 0,$$

and we see that the spectral projection and the Calderón projection of the twisted operator belong to different connected components of the Grassmannian. \square

6. Appendix. Adiabatic Limit of the Cauchy Data Space

We showed that usually the Calderón projection and the Atiyah-Patodi-Singer condition (the spectral projection) belong to different connected components of the Grassmannian. There is, however, a more precise description of the difference between those two boundary conditions. We give a brief review of some results of the beautiful work of Liviu Nicolaescu [18], [19] which are valid in great generality. For explicit calculations characterizing the Cauchy data space $\mathcal{H}_+(\mathcal{D})$ of the Euclidean, untwisted Dirac operator on the 4-ball in terms of the eigenfunctions of the tangential ('spherical') Dirac operator on the 3-sphere see Sudbery, [27], and Sommen, [26].

We have to point out that in his work Nicolaescu considered the case of an odd-dimensional manifold with boundary, but the result holds also in the case of the total Dirac operator \mathcal{D} on an even-dimensional manifold with boundary.

For real, positive R we define the manifold M_R as

$$M_R := ([-R, 0] \times Y) \cup M.$$

The operator \mathcal{D} extends to M_R in a natural way. We study the Calderón projection $\mathcal{P}_+^R(\mathcal{D})$ and the Cauchy data space $\mathcal{H}_+^R(\mathcal{D})$ of the operator \mathcal{D} on M_R . We introduce the notion of the resonance set for the operator \mathcal{D} on M_R :

$$N_R(\mathcal{D}) := \{t \mid H_{<t}(\mathcal{D}) \cap \mathcal{H}_+^R(\mathcal{D}) = \{0\}\}.$$

Here $H_{<t}(\mathcal{D})$ denotes the subspace of $L^2(M; S)$ spanned by eigensections of \mathcal{D} corresponding to eigenvalues smaller than t . Nicolaescu proved that there exists a real number $E(\mathcal{D}) := \sup\{N_R(\mathcal{D})\} \leq 0$. One of the main technical results of his work is the following theorem:

Theorem 6.1 *There exists a positive number a and a Lagrangian subspace \mathcal{L} of $H_{[-a,a]}(\mathcal{D})$ such that:*

$$\lim_{R \rightarrow \infty} \mathcal{H}_+^R(\mathcal{D}) = H_{>a}(\mathcal{D}) \oplus \mathcal{L}. \quad (31)$$

It was pointed out by Nicolaescu that the convergence in the formula (31) is not uniform. There is a more precise result in the so-called non-resonance case.

Definition 6.2 The operator \mathcal{D} is called *non-resonant*, if $E(\mathcal{D}) = \{0\}$.

In this case the convergence is uniform and we have the following result:

Theorem 6.3

$$\lim_{R \rightarrow \infty} \mathcal{H}_+^R(\mathcal{D}) = H_{>0}(\mathcal{D}) \oplus \mathcal{L}^2,$$

where \mathcal{L}^2 denotes the space of the limiting values of so-called extended L^2 -solutions of the operator \mathcal{D} .

Now we can reprove the result of the previous section:

Corollary 6.4 *The Euclidean Dirac operator \mathcal{D}_A on the 4-ball V coupled to any connection A which is pure gauge at the boundary is a resonant operator.*

Proof Assume that the operator \mathcal{D}_A is non-resonant. We know that in our case the tangential operator is invertible, and therefore

$$\lim_{R \rightarrow \infty} \mathcal{H}_+^R(\mathcal{D}_A^+) = H_{>0}(\mathcal{D}_h),$$

which means that the Calderón projection $\mathcal{P}(\mathcal{D}_A^+)$ and the spectral projection $\Pi_{>}(\mathcal{D}_h)$ onto $H_{>0}(\mathcal{D}_h)$ belong to the same connected component of the Grassmannian and hence by [6], Theorem 20.8 $\text{index}(\mathcal{D}_A^+)_{\Pi_{>}} = 0$, but we know that it is equal to $\deg h \neq 0$. \square

7. Addendum. Elementary Proof of the Ninomiya/Tan Result (*sketch, unfinished*)

We offer a second proof of the Ninomiya/Tan result (Proposition 3.5) based on results from [6], where the authors have shown that the index of any elliptic boundary problem for operators of Dirac type is equal to the index of a suitable Toeplitz operator which lives on the boundary (see [6], Theorem 20.1). In fact, instead of applying the Atiyah-Patodi-Singer Index Theorem we apply the general index formula for global elliptic boundary problems of Proposition 1.4 yielding

$$\text{index}(\mathcal{D}_A^+)_{\Pi_{>}} = i(\Pi_{>}, \mathcal{P}_+) = \deg(h).$$

To derive the right equality we recall from (30) the explicit description of the Atiyah-Patodi-Singer projection of the tangential Dirac operator \mathcal{D}_h for the twisted Dirac operator \mathcal{D}_A coupled to a vector potential $A \in \text{Conn}_0(V \times \mathbb{C}^2)$

$$\Pi_{>}(\mathcal{D}_h) = (\text{Id} \otimes h)(\Pi_{>}(\mathcal{D}) \otimes \text{Id})(\text{Id} \otimes h^{-1}). \quad (32)$$

Then the virtual codimension can be written as the index of an elliptic pseudo-differential operator over the closed manifold Y and the Ninomiya/Tan formula follows from the (classical) Atiyah-Singer Index Theorem.

More precisely,

$$\begin{aligned} \text{index}(\mathcal{D}_A^+)_{\Pi_{>}(\mathcal{D}_h)} &= i(\Pi_{>}(\mathcal{D}_h), \mathcal{P}_+(\mathcal{D}_A^+)) = i(\Pi_{>}(\mathcal{D} \otimes_h \text{Id}), \mathcal{P}_+(\mathcal{D} \otimes_A \text{Id})) \\ &= i(\Pi_{>}(\mathcal{D} \otimes_h \text{Id}), \mathcal{P}_+(\mathcal{D} \otimes_d \text{Id})) = i(\Pi_{>}(\mathcal{D} \otimes_h \text{Id}), n\mathcal{P}_+(\mathcal{D})) \\ &= i(\Pi_{>}(\mathcal{D} \otimes_h \text{Id}), \Pi_{>}(\mathcal{D} \otimes_d \text{Id})) + i(\Pi_{>}(\mathcal{D} \otimes_d \text{Id}), n\mathcal{P}_+(\mathcal{D})) \\ &= i(\Pi_{>}(\mathcal{D} \otimes_h \text{Id}), \Pi_{>}(\mathcal{D} \otimes_d \text{Id})) + ni(\Pi_{>}(\mathcal{D}), \mathcal{P}_+(\mathcal{D})). \end{aligned}$$

Note that $\mathcal{P}_+(\mathcal{D} \otimes_d \text{Id}) = \mathcal{P}_+(\mathcal{D}) \otimes \text{Id}_{\mathbb{C}^n} = n\mathcal{P}_+(\mathcal{D})$ and $n = 2$ in our application. The third equality follows from the continuity of taking the Calderón projector: since $\mathcal{D} \otimes_A \text{Id}$ and $\mathcal{D} \otimes_d \text{Id} = n\mathcal{D}$ are connected by a smooth path, the Calderón projectors $\mathcal{P}_+(\mathcal{D} \otimes_A \text{Id})$ and $\mathcal{P}_+(\mathcal{D} \otimes_d \text{Id})$ belong to the same connected component of the Grassmannian and the virtual codimensions coincide.

Now we determine the last sum in the preceding array of equations. The left term does not provide any problems:

$$\begin{aligned} i(\Pi_>(\not{D} \otimes_h \text{Id}), \Pi_>(\not{D} \otimes \text{Id})) &= i((\text{Id} \otimes h)\Pi_>(\not{D} \otimes_d \text{Id})(\text{Id} \otimes h^{-1}), \Pi_>(\not{D} \otimes \text{Id})) \\ &= \deg h \end{aligned}$$

by elementary index theory.

The vanishing of $i(\Pi_>(\not{D}), \mathcal{P}_+(\not{D}))$ is trivial when one uses Kori's result (see Remark 3.4a). For general metrics it follows when one shows that \not{D} on the ball is non-resonant (*must be worked out*).

References

1. Atiyah, M.F.: 1985, 'Topological aspects of anomalies', in: *Anomalies, Geometry, Topology*, Symp. Chicago, Ill., pp. 22-32.
2. Atiyah, M.F., Patodi, V.K., and Singer, I.M.: 1975, 'Spectral asymmetry and Riemannian geometry. I', *Math. Proc. Cambridge Phil. Soc.* **77**, 43-69.
3. Atiyah, M.F., and Singer, I.M.: 1984, 'Dirac operators coupled to vector potentials', *Proc. Nat. Acad. Sci. USA* **81**, 2597-2600.
4. Booß, B., and Bleeker, D.D.: 1985, *Topology and Analysis. The Atiyah-Singer Index Formula and Gauge-Theoretic Physics*, Springer, New York.
5. Booß, B., and Wojciechowski, K.P.: 1989, 'Pseudo-differential projections and the topology of certain spaces of elliptic boundary value problems', *Comm. Math. Phys.* **121**, 1-9.
6. —, —: 1993, *Elliptic Boundary Problems for Dirac Operators*, Birkhäuser, Boston. 1991,
7. Dubrovin, B.A., Fomenko, A.T., and Novikov, S.P.: 1979 *Modern Geometry - Methods and Applications I, II, III*, Nauka, Moscow, (Russian, English translation Springer, Berlin, 1984, 1985, 1990).
8. Fujikawa, K.: 1980, 'Path integral for gauge theories with fermions', *Phys. Rev.* **D21**, 2848-2858.
9. Gilkey, P.B.: 1975, 'The boundary integrand in the formula for the signature and Euler characteristic of a Riemannian manifold with boundary', *Adv. in Math.* **15**, 334-360.
10. —: 1984, *Invariance Theory, the Heat Equation, and the Atiyah-Singer Index Theorem*, Publish or Perish, Wilmington. New revised edition 1995, CRC Press, Boca Raton, Florida.
11. Grubb, G.: 1992, 'Heat operator trace expansions and index for general Atiyah-Patodi-Singer boundary problems', *Commun. Partial Differ. Equations* **17**, 2031-2077.
12. Hitchin, N.: 1974, 'Harmonic spinors', *Adv. Math.* **14**, 1-55.
13. Hortacsu, M., Rothe, K.D., and Schroer, B.: 1980, 'Zero-energy eigenstates for the Dirac boundary problem', *Nucl. Phys.* **B171**, 530-542.
14. Klimek, S., and Wojciechowski, K.P.: 1992, *Adiabatic Cobordism Theorems for Analytic Torsion and η -Invariant*, Preprint, Purdue University, IUPUI, Indianapolis. (To appear in *J. Funct. Anal.*).
15. Kori, T.: 1993, *Index of the Dirac Operator on S^4 and the Infinite Dimensional Grassmannian on S^3* , Preprint, Waseda University, Tokyo.
16. Melrose, R.B., and Piazza, P.: 1993, *Families of Dirac operators, boundaries and the b-calculus*, Preprint.
17. Morchio, G., and Strocchi, F.: 1990, 'Boundary terms, long range effects, and chiral symmetry breaking', in: Mitter, H., and Schweifer, W. (eds), *Fields and Particles*, Proceedings Schladming, Austria, 1990, Springer-Verlag, Berlin-Heidelberg-New York, pp. 171-214.
18. Nicolaescu, L.: 1993, 'The Maslov index, the spectral flow, and splittings of manifolds', *C. R. Acad. Sci. Paris* **317**, Série I, 515-519.
19. —: 1995, 'The Maslov index, the spectral flow, and decompositions of manifolds', (to appear in *Duke Math. J.*).
20. Ninomiya, M., and Tan, C.I.: 1985, 'Axial anomaly and index theorem for manifolds with boundary', *Nucl. Physics.* **257**, 199-225.

21. Palais, R.S. (ed.): 1965, *Seminar on the Atiyah-Singer Index Theorem*, Ann. of Math. Studies 57, Princeton University Press, Princeton.
22. Schmidt, J.R.: 1987, 'Chiral anomaly for the Dirac operator in an instanton background via the Atiyah-Patodi-Singer theorem', *Phys. Rev. D (3)* 36, 2539-2544.
23. Schmidt, J.R., and Bincer, A.M.: 1987, 'Chiral asymmetry and the Atiyah-Patodi-Singer index for the Dirac operator on a four-dimensional ball', *Phys. Rev. D (3)* 35, 3995-4000.
24. Schroer, B.: 1978, 'Topological methods for gauge theories', in: Urban, P. (ed.), *Facts and Prospects of Gauge Theories*, Schladming Lectures 1978, Acta Physica Austriaca, Suppl. XIX, Springer-Verlag, Wien-New York, pp. 155-202.
25. Singer, I.M.: 1988, 'The η -invariant and the index', in: Yau, S.-T. (ed.), *Mathematical Aspects of String Theory*, World Scientific Press, Singapore, pp. 239-258.
26. Sommen, F.: 1981, 'Spherical monogenic functions and analytic functionals on the unit sphere', *Tokyo J. Math.* 4, 427-456.
27. Sudbery, A.: 1979, 'Quaternionic analysis', *Math. Proc. Cambridge Phil. Soc.* 86, 199-225.
28. Vafa, C., and Witten, E.: 1984, 'Eigenvalue inequalities for fermions in gauge theories', *Comm. Math. Phys.* 95, 257-274.
29. Witten, E.: 1988, 'Quantum field theory, Grassmannians, and algebraic curves', *Comm. Math. Phys.* 113, 529-600.
30. Wojciechowski, K.P.: 1983, 'On the spectral flow and the index theorem for flat vector bundles', in: *Partial Differential Equations*, Banach Center Publications, vol. 10, PWN - Polish Scientific Publishers, Warsaw, pp. 413-422.

Liste over tidligere udkomne tekster
tilsendes gerne. Henvendelse herom kan
ske til IMFUFA's sekretariat
tlf. 46 75 77 11 lokal 2263

-
- 217/92 "Two papers on APPLICATIONS AND MODELLING
IN THE MATHEMATICS CURRICULUM"
by: Mogens Niss
- 218/92 "A Three-Square Theorem"
by: Lars Kadison
- 219/92 "RUPNOK - stationær strømning i elastiske rør"
af: Anja Boisen, Karen Birkelund, Mette Olufsen
Vejleder: Jesper Larsen
- 220/92 "Automatisk diagnosticering i digitale kredsløb"
af: Bjørn Christensen, Ole Møller Nielsen
Vejleder: Stig Andur Pedersen
- 221/92 "A BUNDLE VALUED RADON TRANSFORM, WITH
APPLICATIONS TO INVARIANT WAVE EQUATIONS"
by: Thomas P. Branson, Gestur Olafsson and
Henrik Schlichtkrull
- 222/92 On the Representations of some Infinite Dimensional
Groups and Algebras Related to Quantum Physics
by: Johnny T. Ottesen
- 223/92 THE FUNCTIONAL DETERMINANT
by: Thomas P. Branson
- 224/92 UNIVERSAL AC CONDUCTIVITY OF NON-METALLIC SOLIDS AT
LOW TEMPERATURES
by: Jeppe C. Dyre
- 225/92 "HATMODELLEN" Impedansspektroskopi i ultrarent
en-krystallinsk silicium
af: Anja Boisen, Anders Gorm Larsen, Jesper Varmer,
Johannes K. Nielsen, Kit R. Hansen, Peter Baggild
og Thomas Hougaard
Vejleder: Petr Viscor
- 226/92 "METHODS AND MODELS FOR ESTIMATING THE GLOBAL
CIRCULATION OF SELECTED EMISSIONS FROM ENERGY
CONVERSION"
by: Bent Sørensen

- 227/92 "Computersimulering og fysik"
af: Per M.Hansen, Steffen Holm,
Peter Maibom, Mads K. Dall Petersen,
Pernille Postgaard, Thomas B.Schrøder,
Ivar P. Zeck
Vejleder: Peder Voetmann Christiansen
- 228/92 "Teknologi og historie"
Fire artikler af:
Mogens Niss, Jens Høyrup, Ib Thiersen,
Hans Hedal
- 229/92 "Masser af information uden betydning"
En diskussion af informationsteorien
i Tor Nørretranders' "Mærk Verden" og
en skitse til et alternativ baseret
på andenordens kybernetik og semiotik.
af: Søren Brier
- 230/92 "Vinklens tredeling - et klassisk
problem"
et matematisk projekt af
Karen Birkelund, Bjørn Christensen
Vejleder: Johnny Ottesen
- 231A/92 "Elektrondiffusion i silicium - en
matematisk model"
af: Jesper Voetmann, Karen Birkelund,
Mette Olufsen, Ole Møller Nielsen
Vejledere: Johnny Ottesen, H.B.Hansen
- 231B/92 "Elektrondiffusion i silicium - en
matematisk model" Kildetekster
af: Jesper Voetmann, Karen Birkelund,
Mette Olufsen, Ole Møller Nielsen
Vejledere: Johnny Ottesen, H.B.Hansen
- 232/92 "Undersøgelse om den simultane opdagelse
af energiens bevarelse og isærdeles om
de af Mayer, Colding, Joule og Helmholtz
udførte arbejder"
af: L.Arleth, G.I.Dybkjær, M.T.Østergård
Vejleder: Dorthe Posselt
- 233/92 "The effect of age-dependent host
mortality on the dynamics of an endemic
disease and
Instability in an SIR-model with age-
dependent susceptibility
by: Viggo Andreasen
- 234/92 "THE FUNCTIONAL DETERMINANT OF A FOUR-DIMENSIONAL
BOUNDARY VALUE PROBLEM"
by: Thomas P. Branson and Peter B. Gilkey
- 235/92 OVERFLADESTRUKTUR OG POREUDVIKLING AF KOKS
- Modul 3 fysik projekt -
af: Thomas Jessen
-

- 236a/93 INTRODUKTION TIL KVANTE
HALL EFFEKTEN
af: Ania Boisen, Peter Bøggild
Vejleder: Peder Voetmann Christiansen
Erland Brun Hansen
- 236b/93 STRØMSSAMMENBRUD AF KVANTE
HALL EFFEKTEN
af: Anja Boisen, Peter Bøggild
Vejleder: Peder Voetmann Christiansen
Erland Brun Hansen
- 237/93 The Wedderburn principal theorem and
Shukla cohomology
af: Lars Kadison
- 238/93 SEMIOTIK OG SYSTEMEGENSKABER (2)
Vektorbånd og tensorer
af: Peder Voetmann Christiansen
- 239/93 Valgsystemer - Modelbygning og analyse
Matematik 2. modul
af: Charlotte Gjerrild, Jane Hansen,
Maria Hermannsson, Allan Jørgensen,
Ragna Clauson-Kaas, Poul Lützen
Vejleder: Mogens Niss
- 240/93 Patologiske eksempler.
Om sære matematiske fæks betydning for
den matematiske udvikling
af: Claus Dråby, Jørn Skov Hansen, Runa
Ulsøe Johansen, Peter Meibom, Johannes
Kristoffer Nielsen
Vejleder: Mogens Niss
- 241/93 FOTOVOLTAISK STATUSNOTAT 1
af: Bent Sørensen
- 242/93 Brovedligeholdelse - bevar mig vel
Analyse af Vejdirektoratets model for
optimering af broreparationer
af: Linda Kyndlev, Kare Fundal, Kamma
Tulinus, Ivar Zeck
Vejleder: Jesper Larsen
- 243/93 TANKEEKSPERIMENTER I FYSIKKEN
Et 1.modul fysikprojekt
af: Karen Birkelund, Stine Sofia Korremann
Vejleder: Dorthe Posselt
- 244/93 RADONTRANSFORMATIONEN og dens anvendelse
i CT-scanning
Projektrapport
af: Trine Andreasen, Tine Guldager Christiansen,
Nina Skov Hansen og Christine Iversen
Vejledere: Gestur Olafsson og Jesper Larsen
- 245a+b
/93 Time-Of-Flight målinger på krystallinske
halvledere
Specialerapport
af: Linda Szkotak Jensen og Lise Odgaard Gade
Vejledere: Petr Viscor og Niels Boye Olsen
- 246/93 HVERDAGSVIDEN OG MATEMATIK
- LÆREPROCESSER I SKOLEN
af: Lena Lindenskov, Statens Humanistiske
Forskningsråd, RUC, IMFUFA
- 247/93 UNIVERSAL LOW TEMPERATURE AC CON-
DUCTIVITY OF MACROSCOPICALLY
DISORDERED NON-METALS
by: Jeppe C. Dyre
- 248/93 DIRAC OPERATORS AND MANIFOLDS WITH
BOUNDARY
by: B. Booss-Bavnbek, K.P.Wojciechowski
- 249/93 Perspectives on Teichmüller and the
Jahresbericht Addendum to Schappacher,
Scholz, et al.
by: B. Booss-Bavnbek
With comments by W.Abikoff, L.Ahlfors,
J.Cerf, P.J.Davis, W.Fuchs, F.P.Gardiner,
J.Jost, J.-P.Kahane, R.Lohan, L.Lorch,
J.Radkau and T.Söderqvist
- 250/93 EULER OG BOLZANO - MATEMATISK ANALYSE SET I ET
VIDENSKABSTEORETISK PERSPEKTIV
Projektrapport af: Anja Juul, Lone Michelsen,
Tomas Højgård Jensen
Vejleder: Stig Andur Pedersen
- 251/93 Genotypic Proportions in Hybrid Zones
by: Freddy Bugge Christiansen, Viggo Andreasen
and Ebbe Thue Poulsen
- 252/93 MODELLERING AF TILFÆLDIGE FÆNOMENER
Projektrapport af: Birthe Friis, Lisbeth Helmgård,
Kristina Charlotte Jakobsen, Marina Mosbæk
Johannessen, Lotte Ludvigsen, Mette Hass Nielsen
- 253/93 Kuglepakning
Teori og model
af: Lise Arleth, Kåre Fundal, Nils Kruse
Vejleder: Mogens Niss
- 254/93 Regressionsanalyse
Materiale til et statistikkursus
af: Jørgen Larsen
- 255/93 TID & BETINGET UAFHÆNGIGHED
af: Peter Harremoës
- 256/93 Determination of the Frequency Dependent
Bulk Modulus of Liquids Using a Piezo-
electric Spherical Shell (Preprint)
by: T. Christensen and N.B.Olsen
- 257/93 Modellering af dispersion i piezoelektriske
keramikker
af: Pernille Postgaard, Jørnik Rasmussen,
Christina Specht, Mikko Østergård
Vejleder: Tage Christensen
- 258/93 Supplerende kursusedmateriale til
"Lineære strukturer fra algebra og analyse"
af: Mogens Brun Hæsfelt
- 259/93 STUDIES OF AC HOPPING CONDUCTION AT LOW
TEMPERATURES
by: Jeppe C. Dyre
- 260/93 PARTITIONED MANIFOLDS AND INVARIANTS IN
DIMENSIONS 2, 3, AND 4
by: B. Booss-Bavnbek, K.P.Wojciechowski

- 261/93 OPGAVESAMLING
Bredde-kursus i Fysik
Eksamensopgaver fra 1976-93
- 262/93 Separability and the Jones
Polynomial
by: Lars Kadison
- 263/93 Supplerende kursusmateriale til
"Lineære strukturer fra algebra
og analyse" II
af: Mogens Brun Heefelt
- 264/93 FOTOVOLTAISK STATUSNOTAT 2
af: Bent Sørensen
-
- 265/94 SPHERICAL FUNCTIONS ON ORDERED
SYMMETRIC SPACES
To Sigurdur Helgason on his
sixtyfifth birthday
by: Jacques Faraut, Joachim Hilgert
and Gestur Olafsson
- 266/94 Kommensurabilitets-oscillationer i
laterale supergitre
Fysikspeciale af: Anja Boisen,
Peter Bøggild, Karen Birkelund
Vejledere: Rafael Taboryski, Poul Erik
Lindelof, Peder Voetmann Christiansen
- 267/94 Kom til kort med matematik på
Eksperimentarium - Et forslag til en
opstilling
af: Charlotte Gjerrild, Jane Hansen
Vejleder: Bernhelm Booss-Bavnbek
- 268/94 Life is like a sewer ...
Et projekt om modellering af aorta via
en model for strømning i kloakrør
af: Anders Marcussen, Anne C. Nilsson,
Lone Michelsen, Per M. Hansen
Vejleder: Jesper Larsen
- 269/94 Dimensionsanalyse en introduktion
metaprojekt, fysik
af: Tine Guldager Christiansen,
Ken Andersen, Nikolaj Hermann,
Jannik Rasmussen
Vejleder: Jens Højgaard Jensen
- 270/94 THE IMAGE OF THE ENVELOPING ALGEBRA
AND IRREDUCIBILITY OF INDUCED REPRE-
SENTATIONS OF EXPONENTIAL LIE GROUPS
by: Jacob Jacobsen
- 271/94 Matematikken i Fysikken.
Opdaget eller opfundet
NAT-BAS-projekt
vejleder: Jens Højgaard Jensen
- 272/94 Tradition og fornyelse
Det praktiske eleverarbejde i gymnasiets
fysikundervisning, 1907-1988
af: Kristian Hoppe og Jeppe. Guldager
Vejledning: Karin Beyer og Nils Hybel
- 273/94 Model for kort- og mellemdistanceløb
Verifikation af model
af: Lise Fabricius Christensen, Helle Pilemann,
Bettina Sørensen
Vejleder: Mette Olufsen
- 274/94 MODEL 10 - en matematisk model af intravenøse
anæstetikas farmakokinetik
3. modul matematik, forår 1994
af: Trine Andreasen, Bjørn Christensen, Christine
Green, Anja Skjoldborg Hansen. Lisbeth
Helmgaard
Vejledere: Viggo Andreasen & Jesper Larsen
- 275/94 Perspectives on Teichmüller and the Jahresbericht
2nd Edition
by: Bernhelm Booss-Bavnbek
- 276/94 Dispersionsmodellering
Projektrapport 1. modul
af: Gitte Andersen, Rehannah Børup, Lisbeth Friis,
Per Gregersen, Kristina Vejro
Vejleder: Bernhelm Booss-Bavnbek
- 277/94 PROJEKTARBEJDSPÆDAGOGIK - Om tre tolkninger af
problemorienteret projektarbejde
af: Claus Flensted Behrens, Frederik Voetmann
Christiansen, Jørn Skov Hansen, Thomas
Thingstrup
Vejleder: Jens Højgaard Jensen
- 278/94 The Models Underlying the Anaesthesia
Simulator Sophus
by: Mette Olufsen(Math-Tech), Finn Nielsen
(RISØ National Laboratory), Per Føge Jensen
(Herlev University Hospital), Stig Andur
Pedersen (Roskilde University)
- 279/94 Description of a method of measuring the shear
modulus of supercooled liquids and a comparison
of their thermal and mechanical response
functions.
af: Tage Christensen
- 280/94 A Course in Projective Geometry
by Lars Kadison and Matthias T. Kromann
- 281/94 Modellering af Det Cardiovasculære System med
Neural Puls kontrol
Projektrapport udarbejdet af:
Stefan Frello, Runa Ulsøe Johansen,
Michael Poul Curt Hansen, Klaus Dahl Jensen
Vejleder: Viggo Andreasen
- 282/94 Parallelle algoritmer
af: Erwin Dan Nielsen, Jan Danielsen,
Niels Bo Johansen

- 283/94 Grænser for tilfældighed
(en kaotisk talgenerator)
af: Erwin Dan Nielsen og Niels Bo Johansen
- 284/94 Det er ikke til at se det, hvis man ikke
lige ve' det!
Gymnasiematematikens begrundelsesproblem
En specialerapport af Peter Hauge Jensen
og Linda Kyndlev
Vejleder: Mogens Niss
- 285/94 Slow coevolution of a viral pathogen and
its diploid host
by: Viggo Andreasen and
Freddy B. Christiansen
- 286/94 The energy master equation: A low-temperature
approximation to Bässler's random walk model
by: Jeppe C. Dyre
- 287/94 A Statistical Mechanical Approximation for the
Calculation of Time Auto-Correlation Functions
by: Jeppe C. Dyre
- 288/95 PROGRESS IN WIND ENERGY UTILIZATION
by: Bent Sørensen
- 289/95 Universal Time-Dependence of the Mean-Square
Displacement in Extremely Rugged Energy
Landscapes with Equal Minima
by: Jeppe C. Dyre and Jacob Jacobsen
- 290/95 Modellering af uregelmæssige bølger
Et 3.modul matematik projekt
af: Anders Marcussen, Anne Charlotte Nilsson,
Lone Michelsen, Per Mørkegaard Hansen
Vejleder: Jesper Larsen
- 291/95 1st Annual Report from the project
LIFE-CYCLE ANALYSIS OF THE TOTAL DANISH
ENERGY SYSTEM
an example of using methods developed for the
OECD/IEA and the US/EU fuel cycle externality study
by: Bent Sørensen
- 292/95 Fotovoltaisk Statusnotat 3
af: Bent Sørensen
- 293/95 Geometridiskussionen - hvor blev den af?
af: Lotte Ludvigsen & Jens Frandsen
Vejleder: Anders Madsen
- 294/95 Universets udvidelse -
et metaprojekt
Af: Jesper Duelund og Birthe Friis
Vejleder: Ib Lundgaard Rasmussen
- 295/95 A Review of Mathematical Modeling of the
Controlled Cardiovascular System
By: Johnny T. Ottesen
- 296/95 RETIKULER den klassiske mekanik
af: Peder Voetmann Christiansen
- 297/95 A fluid-dynamical model of the aorta with
bifurcations
by: Mette Olufsen and Johnny Ottesen
- 298/95 Mordet på Schrödingers kat - et metaprojekt om
to fortolkninger af kvantemekanikken
af: Maria Hermannsson, Sebastian Horst,
Christina Specht
Vejledere: Jeppe Dyre og Peder Voetmann Christiansen
- 299/95 ADAM under figenbladet - et kig på en samfunds-
videnskabelig matematisk model
Et matematisk modelprojekt
af: Claus Dræby, Michael Hansen, Tomas Højgård Jensen
Vejleder: Jørgen Larsen
- 300/95 Scenarios for Greenhouse Warming Mitigation
by: Bent Sørensen
- 301/95 TOK Modellering af træers vækst under påvirkning
af ozon
af: Glenn Møller-Holst, Marina Johannessen, Birthe
Nielsen og Bettina Sørensen
Vejleder: Jesper Larsen
- 302/95 KOMPRESSORER - Analyse af en matematisk model for
aksialkompressorer
Projektrapport af: Stine Bøggild, Jakob Hilmer,
Pernille Postgaard
Vejleder: Viggo Andreasen
- 303/95 Masterlignings-modeller af Glasovergangen
Termisk-Mekanisk Relaksation
Specialerapport udarbejdet af:
Johannes K. Nielsen, Klaus Dahl Jensen
Vejledere: Jeppe C. Dyre, Jørgen Larsen
- 304a/95 STATISTIKNOTER Simple binomialfordelingsmodeller
af: Jørgen Larsen
- 304b/95 STATISTIKNOTER Simple normalfordelingsmodeller
af: Jørgen Larsen
- 304c/95 STATISTIKNOTER Simple Poissonfordelingsmodeller
af: Jørgen Larsen
- 304d/95 STATISTIKNOTER Simple multinomialfordelingsmodeller
af: Jørgen Larsen
- 304e/95 STATISTIKNOTER Mindre matematisk-statistisk opslagsværk
indeholdende bl.a. ordforklaringer, resuméer og
tabeller
af: Jørgen Larsen

- 305/95 The Maslov Index:
A Functional Analytical Definition
And The Spectral Flow Formula

By: B. Booss-Bavnbek, K. Furutani
- 306/95 Goals of mathematics teaching

Preprint of a chapter for the forthcoming International Handbook of Mathematics Education (Alan J. Bishop, ed)

By: Mogens Niss
- 307/95 Habit Formation and the Thirdness of Signs

Presented at the semiotic symposium

The Emergence of Codes and Intensions as a Basis of Sign Processes

By: Peder Voetmann Christiansen
- 308/95 Metaforer i Fysikken

af: Marianne Wilcken Bjerregaard,
Frederik Voetmann Christiansen,
Jørn Skov Hansen, Klaus Dahl Jensen
Ole Schmidt

Vejledere: Peder Voetmann Christiansen og
Petr Viscor
- 309/95 Tiden og Tanken

En undersøgelse af begrebsverdenen Matematik
udført ved hjælp af en analogi med tid

af: Anita Stark og Randi Petersen

Vejleder: Bernhelm Booss-Bavnbek
- 310/96 Kursusmateriale til "Lineære strukturer fra
algebra og analyse" (E1)

af: Mogens Brun Heefelt
- 311/96 2nd Annual Report from the project

LIFE-CYCLE ANALYSIS OF THE TOTAL DANISH
ENERGY SYSTEM

by: Hélène Connor-Lajambe, Bernd Kuemmel,
Stefan Krüger Nielsen, Bent Sørensen